

***ald* of *Mycobacterium tuberculosis* Encodes both the Alanine Dehydrogenase and the Putative Glycine Dehydrogenase**

Michelle M. Giffin, Lucia Modesti, Ronald W. Raab,
Lawrence G. Wayne and Charles D. Sohaskey
J. Bacteriol. 2012, 194(5):1045. DOI: 10.1128/JB.05914-11.
Published Ahead of Print 30 December 2011.

Updated information and services can be found at:
<http://jb.asm.org/content/194/5/1045>

SUPPLEMENTAL MATERIAL

These include:

[Supplemental material](#)

REFERENCES

This article cites 60 articles, 34 of which can be accessed free
at: <http://jb.asm.org/content/194/5/1045#ref-list-1>

CONTENT ALERTS

Receive: RSS Feeds, eTOCs, free email alerts (when new
articles cite this article), [more»](#)

Information about commercial reprint orders: <http://journals.asm.org/site/misc/reprints.xhtml>
To subscribe to to another ASM Journal go to: <http://journals.asm.org/site/subscriptions/>

ald of *Mycobacterium tuberculosis* Encodes both the Alanine Dehydrogenase and the Putative Glycine Dehydrogenase

Michelle M. Giffin,^a Lucia Modesti,^a Ronald W. Raab,^b Lawrence G. Wayne,^a and Charles D. Sohaskey^a

Department of Veterans Affairs Medical Center, Long Beach, California, USA,^a and Department of Integrated Science and Technology, James Madison University, Harrisonburg, Virginia, USA^b

The putative glycine dehydrogenase of *Mycobacterium tuberculosis* catalyzes the reductive amination of glyoxylate to glycine but not the reverse reaction. The enzyme was purified and identified as the previously characterized alanine dehydrogenase. The *Ald* enzyme was expressed in *Escherichia coli* and had both pyruvate and glyoxylate aminating activities. The gene, *ald*, was inactivated in *M. tuberculosis*, which resulted in the loss of all activities. Both enzyme activities were found associated with the cell and were not detected in the extracellular filtrate. By using an anti-Ald antibody, the protein was localized to the cell membrane, with a smaller fraction in the cytosol. None was detected in the extracellular medium. The *ald* knockout strain grew without alanine or glycine and was able to utilize glycine but not alanine as a nitrogen source. Transcription of *ald* was induced when alanine was the sole nitrogen source, and higher levels of Ald enzyme were measured. Ald is proposed to have several functions, including ammonium incorporation and alanine breakdown.

Mycobacterium tuberculosis is the causative agent of tuberculosis, and one of the most successful human pathogens. It was responsible for approximately 2 million deaths in 2008, while currently almost one-third of the world's population is infected with this organism. Research with *M. tuberculosis* has described a pathogen uniquely adapted to the wide range of harsh environments presented by the host. Much of this work has focused on the microbe's metabolism, with the idea of identification of novel enzymes or pathways to target for drug development.

One of these environmental factors is nitrogen availability. Very little is known about the nitrogen sources used by *M. tuberculosis* *in vivo*. *M. tuberculosis* can utilize many amino acids for nitrogen, including alanine and glycine (29). Mutants of *M. tuberculosis* unable to synthesize proline retained the ability to replicate in the human macrophage cell line THP-1 (35), while other amino acid auxotroph mutants were attenuated (3, 17, 22, 52). A *Mycobacterium bovis* BCG mutant unable to make methionine showed survival in mice similar to the wild-type strain (32). This suggests some amino acids are available *in vivo* and serve as nutrients for *M. tuberculosis*.

The enzyme glycine dehydrogenase was first described in *M. tuberculosis* in 1962 (16). This enzyme was detected by the reductive amination of glyoxylate to glycine concurrent with the oxidation of NADH to NAD⁺ (Fig. 1). This reaction represents glyoxylate reductive aminase (GxRA) activity. The activity corresponding to the reverse reaction, catalyzed by glycine dehydrogenase (GDH), was not detected. The expression of glyoxylate reductive amination by a putative glycine dehydrogenase in *M. tuberculosis* has been characterized in nonreplicating persistent (NRP) cultures (58). In the Wayne model of dormancy, sealed cultures of *M. tuberculosis* create a microaerobic environment (NRP-1), which subsequently develops into the anaerobic stage (NRP-2) (58). GxRA activity was induced during microaerobic NRP-1, with the strongest activity in anaerobic NRP-2 cultures. It was proposed that the role of this enzyme was to maintain redox balance by recycling NADH/NAD⁺ during interruption of aerobic respiration (59, 60).

The naming of the glycine dehydrogenase was based on the

similarity of the glyoxylate reductive aminase reaction to that catalyzed by L-alanine dehydrogenase (Ald; L-alanine:NAD⁺ oxidoreductase; EC 1.4.1.1) (15). Alanine dehydrogenase catalyzed the reductive amination of pyruvate to L-alanine (PvRA), but the reverse reaction, the oxidative dehydrogenation of L-alanine (ALD), was also detected. Alanine dehydrogenases are well-studied enzymes found in a wide range of bacterial species. In mycobacteria, it was first identified as an enzyme absent from the vaccine strains of *M. bovis* BCG but present in virulent *M. tuberculosis* (2). It was suggested that impairment of *M. bovis* BCG replication in humans due to the lack of a functional alanine dehydrogenase inhibited the development of protective immunity (44).

There have been several attempts to identify the gene encoding the putative glycine dehydrogenase. *gcvB* (Rv1832), annotated in the *M. tuberculosis* genome as a glycine dehydrogenase gene, most likely encodes the P protein of the glycine cleavage system (60). In *Mycobacterium smegmatis* pyruvate and glyoxylate aminase activities comigrated on a native polyacrylamide gel (53), suggesting one enzyme for both activities. However, a knockout mutant of the alanine dehydrogenase gene *ald* in *M. smegmatis* lost alanine dehydrogenase activity but retained glycine dehydrogenase activity (14). Moreover, *M. bovis* does not produce alanine dehydrogenase, but glycine dehydrogenase activity has been reported (4, 27). Thus, the identity of this unique enzyme is unknown.

In this study *ald* was shown to encode both alanine dehydrogenase and glyoxylate reductive aminase (glycine dehydrogenase) activities. This was determined by both biochemical and genetic methods. This dual function enzyme was localized to the cell

Received 1 August 2011 Accepted 20 December 2011

Published ahead of print 30 December 2011

Address correspondence to Charles Sohaskey, Chuck@Sohaskey.com.

Supplemental material for this article may be found at <http://jb.asm.org/>.

Copyright © 2012, American Society for Microbiology. All Rights Reserved.

doi:10.1128/JB.05914-11




FIG 1 Possible reactions of alanine dehydrogenase. GDH activity was not detected.

membrane and cytosol. It plays an essential role in the utilization of alanine, but not glycine, as a nitrogen source.

MATERIALS AND METHODS

Strains and media. *M. tuberculosis* H37Rv and Erdman strains were from the culture collection of this laboratory. Mycobacterial cultures were grown at 37°C in Dubos Tween-albumin broth (DTA; Difco, Detroit, MI). For studies with nitrogen or carbon sources, minimal lysis-inducing medium (LIM) was used with amino acids at 10 mM (45). Aerobic cultures were incubated on a model G24 rotary shaker-incubator (New Brunswick Scientific, Edison, NJ). For hypoxic cultures (Wayne model), slow magnetic stirring in sealed tubes with a headspace ratio of 0.5 were prepared as previously described (58).

Hygromycin was used at a concentration of 50 µg/ml, and gentamicin was used at 10 µg/ml. All antibiotics and chemicals were from Sigma (St. Louis, MO).

Purification of glycine dehydrogenase. *M. tuberculosis* was grown in flasks of 400 ml of DTA in the Wayne model (58). Cells were harvested by centrifugation, washed twice with 100 mM phosphate buffer (pH 7.0), and disrupted by sonication (48). DEAE-Sephacel was used to remove excessive lipids. The gel was first equilibrated with 10 mM phosphate buffer (pH 7.0), then the cell extracts were loaded, and finally protein was eluted with 200 mM phosphate buffer (pH 7.0) (16). Glycine dehydrogenase was further purified at Athena Environmental Sciences, Inc. (Baltimore, MD) by using a two-step chromatographic fractionation procedure. In the first step, hydrophobic interaction chromatography was used. Cell extract was fractionated on a phenyl-Sepharose HiTrap column (Amersham Pharmacia Biotech, Piscataway, NJ) at pH 7.0. Next, anion-exchange chromatography was used. Samples with glyoxylate reductive aminase activity were fractionated on a 1-ml Mono Q HR5/5 column (Amersham Pharmacia Biotech). The fraction with the greatest activity was concentrated by ultrafiltration with a Microcon YM-3 column (Millipore, Billerica, MA) and washed three times with 10 mM ammonium acetate (pH 6.8). The sample was subjected to tryptic digestion, fractionated by microcapillary reverse-phase high-performance liquid chromatography, and then run through a nano-electrospray ionization source of an ion trap mass spectrometer. The tandem mass spectrometry data were correlated with known sequences by using the Sequest and HMF computer algorithms (13).

Preparation of cell extracts for enzyme studies. *M. tuberculosis* was grown aerobically to either mid-log phase with an approximate optical density at 580 nm (OD_{580}) of 0.4 or to NRP-2 after 250 h in the Wayne model. To harvest, cell suspensions were placed on ice for 2 h and then washed twice by centrifugation with cold 100 mM phosphate buffer (pH

7.0) containing 0.05% Tween 80. Cells were disrupted using a minibeat-beater (BioSpec Products, Bartlesville, OK) with 6 repetitions of 30-s bursts and with cooling on ice between each burst. Extracts were clarified by centrifugation and then sterilized by filtration through a 0.22-µm-pore-size SpinX filter (Corning, Corning, NY). The protein concentration was determined with the Bio-Rad (Hercules, CA) protein assay reagents.

Assay of enzyme activity. The assay for GxRA activity for the identification of the putative glycine dehydrogenase was based on the original protocol for measuring the rate of oxidation of NADH during the reductive amination of glyoxylate (16). Approximately 50 µg of cell extract protein or 1 µg of His-tagged Ald was added to a 1-ml UV-transparent cuvette containing 100 mM phosphate buffer. Ammonium sulfate to 400 mM and sodium glyoxylate (pH 6.4) to 50 mM were added. The reactions were started with the addition of NADH to 80 µM and measured by the rate of decrease of the A_{340} on a Libra S32PC spectrophotometer (Biochrome, Cambridge, United Kingdom). A control reaction without ammonium sulfate was included to measure the background, which was subtracted from the experimental values. Glutamine, when employed, was used at 40 mM.

Reductive amination of pyruvate to alanine (PvRA) was measured in an identical manner to that of GxRA activity, except that sodium pyruvate was used in place of glyoxylate. In some experiments pyruvate was replaced with α -ketoglutaric acid, lithium hydroxypyruvate, sodium glycolate, sodium acetate, sodium oxaloacetate, or methylglyoxal at a final concentration of 20 mM.

For the oxidative deamination reaction (ALD) in which alanine was converted to pyruvate, L-alanine at 100 mM and NAD^+ at 1 mM were used (15). Glycine, in place of alanine, was used to detect GDH activity. In some experiments alanine was replaced with aspartate, glutamate, glycinate, proline, or serine at 40 mM.

To determine the pH profile, 4 different buffers were used with overlapping pH ranges: 100 mM phosphate buffer (pH 4.5 to 9.0), 100 mM citrate buffer (pH 3.0 to 6.0), 100 mM Tris-HCl buffer (pH 7.0 to 9.0), and 100 mM carbonate buffer (pH 8.0 to 11.0). Other components were at the same concentrations listed above. All assays were performed in triplicate at room temperature.

To determine the K_m values, the reductive amination reaction or oxidative deamination reaction was performed in 100 mM phosphate buffer at pH 8.5. The concentrations of pyruvate, glyoxylate, ammonium, alanine, NADH, and NAD^+ were varied at least 10-fold above and below the apparent K_m . Other substrates were at the standard concentrations previously mentioned. K_m and V_{max} values were determined from a Lineweaver-Burk plot.

For all reactions, activities were determined from the linear sections of curves. An extinction coefficient of 6,300 $M^{-1} cm^{-1}$ was used. Specific activity was expressed as µmoles of NADH oxidized per min per mg of cell extract protein. The pH was measured after each reaction was complete. All reactions were performed at room temperature.

Three-dimensional crystal structures were analyzed for substrate specificity by using the program COOT (12).

Cloning and expression of histidine-tagged alanine dehydrogenase. The *ald* gene was amplified from *M. tuberculosis* H37Rv by PCR using the primers AscI and NotI (Table 1). The fragment produced was digested with NotI and AscI, ligated into pET DUET (New England BioLabs, Ipswich, MA), and electroporated into *Escherichia coli* ER2566. The resulting plasmid was named pAld-His1.

For expression of the histidine-tagged alanine dehydrogenase, 1 liter of Luria broth with ampicillin was inoculated with ER2566(pAld-His1) and grown with shaking at 37°C to an OD_{600} of between 0.5 and 0.8. Isopropyl- β -D-thiogalactopyranoside (IPTG) was added to 500 µM, and incubation was continued at room temperature for an additional 4 h. Cells were harvested by centrifugation and resuspended in 30 ml of lysis buffer (50 mM NaH_2PO_4 , 300 mM NaCl, 10 mM imidazole) with 1 mg/ml of lysozyme. Cells were disrupted by sonication with 3 repeats for 15 s. The sonicate was cleared by centrifugation, and the His-tagged Ald purified

TABLE 1 Oligonucleotide primers used in this study

Name	Sequence (5'→3') ^a	Purpose in this study ^b
p58	ACTCGAGTGGCGAACGGGTGAGTAACACGT	Q-PCR of 16S
p59	AGGCCGTCACCCACCAACAAGCTGATAGG	Q-PCR of 16S
p169	GCCGCGTCACGTGCTTGACCCGATGCGTTG	Q-PCR of <i>ald</i>
p170	AACCTGGGCGCGAGTCGACCGGCGACTTC	Q-PCR of <i>ald</i>
p171	CCATGCAGCAAGCTTAGGATTCTGCGGTCC	Cloning of <i>ald</i>
p172	GCCACTACCGGTCTATCGGAGCCAATCTG	Cloning of <i>ald</i>
p189	AAGATCAGGGACGCCCGGTACCTGCGACG	Cloning of <i>ald</i>
p190	GAACAAACCTTTGCCCGGATCGGCGCACC	Cloning of <i>ald</i>
AscI	GTGTGGCGCGCCTAATGCGCGTCGGTATT CCGACCGAG	His-Ald
NotI	AAGGAAAAAGCGGCCGCTCAGGCCAGCAC TCGGGCGGCTCGG	His-Ald

^a Mismatches used to create restriction sites for *ald* cloning are indicated by underlining.

^b Q-PCR, quantitative reverse transcriptase PCR. His-Ald, PCR to create the His-tagged *ald* expression plasmid.

was loaded onto a Qiagen 1.5-ml Ni-resin column (Qiagen, Valencia, CA). The column was washed three times with wash buffer (50 mM NaH₂PO₄, 300 mM NaCl, 20 mM imidazole) and then eluted with 4 ml of elution buffer (50 mM NaH₂PO₄, 300 mM NaCl, 250 mM imidazole) as per the manufacturer's instructions. The purified protein was dialyzed twice against 3 liters of phosphate-buffered saline and frozen at -80°C until analyzed (see Fig. S1 in the supplemental material).

Construction of the *ald* knockout. The *ald* gene of *M. tuberculosis* H37Rv was amplified from genomic DNA with primers p171 and p172 (Table 1). The fragment was cloned with the TOPO Blunt cloning kit (Invitrogen, Carlsbad, CA). The *ald* gene was cloned into the suicide vector pJQ200KS (38), followed by the insertion of *hyg*, a hygromycin resistance marker, into the BamHI and XbaI sites in *ald*. A 53-bp fragment of *ald* DNA was deleted in the process. The plasmid was electroporated into *M. tuberculosis* followed by selection on DTA agar plates with 25 µg/ml hygromycin. Inserts of the plasmid into the chromosomal *ald* were identified by Southern analysis (Fig. 2). To create the double-crossover knockout mutant RVW7, clones were replated on DTA agar containing 2% sucrose and hygromycin. Southern analysis was used to verify the correct structure. The wild-type strain produced a fragment of 3.8 kb, while the knockout strain RVW7 produced a larger fragment of 4.8 kb.

To create the complementing plasmid, *ald* was amplified with primers p189 and p190 (Table 1) and cloned into the insertional plasmid pUC-GEN-INT (30). After the *ald* sequence was verified, the plasmid was electroporated into RVW7. Selection was for both gentamicin and hygromycin resistance. Southern analysis was used to verify the presence of the

inactive *ald* gene (band size, 4.8 kb), as well as the insertion of the second complementing *ald* of pAld-Gen21 (band size, 8.8 kb) (Fig. 2).

Digoxigenin-labeled probes for Southern analysis were created by PCR (Roche Diagnostics, Indianapolis, IN) using p169 and p170 as primers.

Subcellular localization of Ald. To determine the location of Ald, *M. tuberculosis* was grown in modified DTA without bovine serum albumin and with Tween 80 replaced by Tyloxapol. Aerobic cultures were grown to mid-log phase (OD₅₈₀ ~0.4), and NRP-2 cultures were grown for 300 h in the Wayne model. Cells were harvested by centrifugation, resuspended in 100 mM phosphate buffer (pH 7.2), and disrupted by bead beating. Spent medium was filter sterilized and concentrated over a Centrplus column (Millipore). Phenylmethanesulfonyl fluoride was added to all fractions at 44 µg/ml.

Eighty micrograms of protein from each sample was separated on a 10% Tris-HEPES IGels Long-Life gel (NuSep, Lane Cove, NSW, Australia). HBT-10, the mouse anti-Ald antibody, was the primary antibody (28). The secondary antibody for experiments with culture filtrate was goat anti-mouse IgG (Fab-specific)-alkaline phosphatase-conjugated antibody (Sigma, St. Louis, MO), and detection was with 5-bromo-4-chloro-3-indolylphosphate-nitroblue tetrazolium (Amresco, Solon, OH).

The cell extract was separated into different fractions by centrifugation (19, 33). Cell extracts were centrifuged at 27,000 × g at 4°C for 30 min to pellet the cell wall. The supernatant was centrifuged at 100,000 × g at 4°C for 2 h to pellet the cell membrane components. Both of these fractions were washed with phosphate-buffered saline. The final supernatant was taken as the soluble cytosol fraction. Ten micrograms of protein from each sample was separated by SDS-PAGE and probed with HBT-10. The secondary antibody was goat anti-mouse IgG (Fab-specific)-horseradish phosphatase-conjugated antibody (Sigma, St. Louis, MO). Detection was performed with the Pierce enhanced chemiluminescence Western blotting substrate (Thermo, Rockford, IL).

RNA isolation. Cultures were quickly harvested by centrifugation, and the cell pellet was resuspended in 1 ml of RNAwiz (Ambion, Austin, TX) before being disrupted in a minibeatbeater by three 1-min pulses. Samples were centrifuged for 5 min at 10,000 × g, and the supernatants was extracted once with CHCl₃ followed by precipitation of the RNA with isopropanol. DNA was removed by treatment with DNase (Roche, Indianapolis, IN) for 2 h at 37°C. RNA was purified with the RiboPure bacteria kit (Ambion) and treated for a second time with DNase, which was then inactivated at 70°C for 5 min.

Quantitation of mRNA levels. cDNA was produced with the RETROscript kit (Ambion) according to the manufacturer's instructions. Approximately 150 ng of RNA was added to a mixture of antisense primers at 1 µM. Duplicate samples without Moloney murine leukemia virus reverse transcriptase were prepared as negative controls.

Real-time quantitative PCR was performed with the Brilliant SYBR green QPCR master mix kit (Stratagene, La Jolla, CA). Reactions were




FIG 2 Southern blot analyses of the *ald* insertional mutant and complement. (A) DNA was isolated from each strain, cut with SacI, and analyzed with a probe specific for *ald*. RVW7, Δ *ald*::*hyg*; Ald21, RVW7(pAld-Gen21). The approximate sizes are indicated on the left. (B) Gene organization of *ald*, showing the insertion of *hyg*. Arrows indicate open readings frames and the direction of transcription. X, XbaI site; B, BamHI site; S, SacI site. *ald'* is the disrupted *ald*. The *ald* probe binding site is shown.

TABLE 2 Proteins identified in a purified GxRA sample from a hypoxic NRP culture^a

Gene	Protein name	Protein description
Rv0350	DnaK	Heat shock 70-kDa protein
Rv2031c	HspX	α -Crystallin/URB-1
Rv2032	Acg	Unknown function
Rv2780	Ald	L-Alanine dehydrogenase
Rv2626c	Hrp1	Hypoxic response protein 1
Rv1133c	MetE	Methyltetrahydropteroyltrimethylglutamate-homocysteine methyltransferase
Rv1070c	EchA8	Enoyl-CoA hydratase/isomerase family protein

^a Proteins were identified by mass spectrometry after a two-step chromatographic purification for glycine dehydrogenase.

performed in a volume of 20 μ l containing two gene-specific primers (Table 1) at 0.1 μ M, 25 μ l of 2 \times master mix, and 5 μ l of cDNA. A 10-fold dilution series of RNA was created, and the 16S RNA was measured to create a standard curve. 16S RNA is stable during long-term persistence in the Wayne model (11). Rv0239 was included as a second standard (21). Amplification was performed in the ICycler (Bio-Rad, Hercules, CA) with sampling during elongation. An initial denaturation of 10 min at 95°C was performed, followed by 30 cycles of 95°C for 30 s and 68°C for 1 min. All samples were run on a 2% agarose gel to verify that only a single band was produced. Each gene was analyzed from 3 independent RNA samples.

RESULTS

Isolation of the putative glycine dehydrogenase. The putative glycine dehydrogenase was purified from *M. tuberculosis* H37Rv by two-step purification. First hydrophobic interaction and then anion-exchange chromatography were used. Seven proteins were isolated and identified, including some of unknown function (Table 2). One enzyme with a known function was L-alanine dehydrogenase (Rv2780), which catalyzes the reductive amination of pyruvate to L-alanine with the oxidation of NADH (Fig. 1). It also catalyzes the reverse reaction: the oxidative deamination reaction of alanine to pyruvate with the reduction of NAD⁺ (15). The isolation of alanine dehydrogenase was not unexpected, as the procedure to isolate glycine dehydrogenase was based, in part, on the purification of NADH binding proteins.

Inactivation of *ald*. To determine if the Ald enzyme was also the putative glycine dehydrogenase, two approaches were used. First, the gene was inactivated in *M. tuberculosis*, and second, in order to biochemically reconstitute the enzyme, it was expressed and purified from *E. coli*. The gene encoding the alanine dehydrogenase, *ald*, is located between two genes transcribed in the opposite direction, suggesting it is a monocistronic gene. If *ald* encodes both the alanine and putative glycine dehydrogenases, then loss of this gene should result in the loss of these activities. To test this, a hygromycin marker was inserted into *M. tuberculosis*, inactivating *ald* and creating the strain RVW7 (Fig. 2).

Glycine dehydrogenase has been detected in hypoxic cultures, and alanine dehydrogenase is generally detected in aerobic cultures (2, 24, 58). Both PvRA and GxRA activities were measured in cell extracts from aerobically growing and from nonreplicating persistent cultures of the Wayne model (Fig. 3). Low PvRA and GxRA activities were detected in aerobic cultures. *M. tuberculosis* from anaerobic cultures showed a 2-fold induction of PvRA activity and 9-fold induction of GxRA activity (Fig. 3B). RVW7 showed neither type of activity under either condition.

To complement the knockout mutant, *ald* plus 527 bp of up-

stream DNA was cloned into the insertional vector pUC-GM-INT (30) to make pAld-Gen21. This plasmid was electroporated into RVW7, creating RVW7(pAld21) (Fig. 2). This strain was tested for GxRA and PvRA activities in both aerobic and NRP cultures (Fig. 3). Both were similar to the wild type (WT), suggesting that insertional inactivation of *ald* was responsible for the loss of PvRA and GxRA activities, which could be complemented with a functional copy of *ald*.

Enzymatic activity of Ald. The alanine dehydrogenase of *M. tuberculosis* was expressed as a histidine-tagged protein and purified from *E. coli*. The enzymatic activity of the His-tagged protein was first verified by determining the pH dependence for the alanine dehydrogenase reactions (Fig. 4A). For the PvRA activity, the optimal pH was 8.5, similar to the previously reported ranges of 8.5 to 9 (15) and 7.0 to 7.5 (24). For the ALD activity, the optimal pH was 9.0, lower than the reported range of 9.5 to 11 (15, 24).

The His-tagged Ald was analyzed for the glyoxylate reductive aminating activity (GxRA) attributable to glycine dehydrogenase. This activity was detected, but GDH activity was not (Fig. 4B). Glutamate was tested for its ability to replace ammonium sulfate, but only weak activity was seen. Therefore, the alanine dehydrogenase enzyme of *M. tuberculosis* also has the glyoxylate aminase activity, which defines the putative glycine dehydrogenase (16). To compare the enzyme activities, GxRA activity at different pH values was measured. This activity was detected with an optimal pH of 8.5 (Fig. 4B), higher than the reported optimum pH of 6.2 (16). These results, taken together with those from the knockout mutant of *M. tuberculosis*, prove that *ald* encodes an enzyme having both PvRA and GxRA activities.

To further characterize this enzyme, the ability to utilize other substrates was tested. In a reductive amination reaction, there was no activity with 2-ketoglutarate, malate, acetate, or glycolate (Fig. 5A). Approximately 60% activity was seen with hydroxypyruvate, but only if ammonium sulfate was included in the reaction mixture, indicating Ald does not have hydroxypyruvate reductase activity. Low activity was detected with oxaloacetate and methylglyoxal. For the oxidative deamination reaction, no activity was seen with proline, aspartate, or serine (Fig. 5B).

The crystal structure of the *M. tuberculosis* alanine dehydrogenase indicates that Arg15 and Lys75 stabilize pyruvate in the active site by binding to the carboxyl group by hydrogen bonds (1, 51). The conserved residues His96 and Asp270 are the catalytic groups. Computer modeling suggests that glyoxylate, hydroxypyruvate, and oxaloacetate, but not α -ketoglutarate, fit into the active site of the holo enzyme (data not shown). However, as the sizes of the substrate increased, they possibly put increasing stress on the active site and the stabilizing amino acids, which resulted in a decline in activity (Fig. 5B). Alanine, glycine, and serine also fit into the active site of the holo but not the apo conformation, based on computer modeling. Why alanine serves as a substrate but glycine or serine do not is not clear.

Enzyme kinetics. The K_m values of His-Ald were determined (Table 3). For pyruvate, the V_{max} was 5.8 mmol/min/mg of protein, and for glyoxylate it was 2.5 mmol/min/mg of protein. The apparent K_m of Ald for pyruvate was 2.8 mM, which was previously reported to range from 0.76 to 1.45 mM (1, 24). The apparent K_m for glyoxylate was 5.5 mM, close to the published range of 0.22 to 5.4 mM (16, 59). The ranges for other apparent K_m values are the following: ammonium, 35 to 2,900 mM; NADH, 9 to 98




FIG 3 (A and B) NADH breakdown with glyoxylate in aerobic cultures (A) or nonreplicating persistent stage 2 cultures (B). Filled symbols and solid lines show data for complete reactions, while empty symbols and dotted lines are for reactions without ammonium sulfate. Circles, WT; triangles, RVW7; squares, RVW7(pAld-Gen21). The standard deviations are shown. (C and D) PvRA and GxRA activities of the Δald mutant in aerobic cultures (C) or nonreplicating persistent stage 2 cultures (D). PvRA is involved in the reductive amination of pyruvate to alanine. GxRA is involved in the reductive amination of glyoxylate to glycine. RVW7, Δald ; Ald21, RVW7(pAld-Gen21). Specific activity is reported in μ moles of NADH oxidized/min/ μ g of protein. The standard deviations are shown.

μ M; alanine, 14 to 16 mM; NAD^+ , 310 μ M (1, 16, 16, 24, 59). The differences may be due to buffer composition and pH, which can affect substrates. Ammonium concentrations, for example, are affected by pH. At higher pHs the level of NH_4^+ , the actual sub-

strate, is higher. Here, a pH of 8.5 in phosphate buffer was used, which is higher than that reported for other studies.

Subcellular localization of Ald. Alanine dehydrogenase is reported to be a secreted enzyme, with high levels found in the




FIG 4 Activity of His-tagged Ald at different pH levels. (A) Pyruvate reductive amination activity (circles) and alanine dehydrogenase activity (squares). (B) Glyoxylate reductive amination activity (circles), glycine dehydrogenase activity (squares), and glyoxylate reductive amination activity with glutamine in place of ammonium sulfate (triangles). Specific activity is shown in μ moles of NADH oxidized/min/ μ g of protein. The standard deviations are shown.




FIG 5 Activity of the Ald enzyme with alternate substrates. (A) Reductive amination reaction with pyruvate (Pyv), glyoxylate (Glx), α -ketoglutarate (aKt), hydroxypyruvate (Hpv), malate (Mal), acetate (Ace), glycolate (Glc), oxaloacetate (Oaa), and methylglyoxal (Mg). (B) Oxidative deamination reaction with alanine (Ala), glutamate (Glu), proline (Pro), aspartate (Asp), serine (Ser), and glycine (Gly). The specific activity of the His-tagged Ald is reported in $\mu\text{moles of NADH oxidized/min}/\mu\text{g of protein}$. The standard deviations are shown.

medium (2, 39, 41), while glyoxylate reductive aminase activity is associated with cell extracts (58). The location of Ald for *M. tuberculosis* was examined in aerobic and NRP cultures by two methods. For the first method, enzyme activity was measured, and for the second an Ald-specific antibody was used. *M. tuberculosis* H37Rv was grown to either mid-log phase in aerobic cultures or to NRP-2. The cells were concentrated by centrifugation, while the cell-free spent medium was removed and concentrated. Packed cells were disrupted, and both PvRA and GxRA activities were determined for each fraction (Table 4). Neither activity was detected in culture filtrates from aerobic or NRP cultures. Strong activity was measured in NRP cell lysates, with lower levels from the aerobic cell lysates. Differences in Ald location related to virulence have been reported for different strains (2, 39), and so the experiment was repeated with the more virulent Erdman strain (Table 4). Activity was detected only in cell lysates and not in culture filtrates.

It was possible the Ald enzyme excreted from the cell was inactive. Therefore, the protein was detected with an anti-Ald antibody (28). Again, alanine dehydrogenase was detected only in the cell extract from NRP cultures (Fig. 6A). Cell extracts were separated into cell wall, membrane, and soluble fractions. Equal amounts of protein were separated by SDS-PAGE and probed with the anti-Ald antibody. Most of the protein (90%) was associated with the cell membrane, with a smaller amount (10%) in the soluble fraction (Fig. 6B).

Role of Ald in utilization of amino acids. Ald, functioning as an alanine dehydrogenase, is essential for the growth of *M. bovis* when alanine is the sole nitrogen source (9), but the role of GxRA activity in the synthesis or breakdown of glycine is not known. The roles of the various enzymatic activities of Ald in *M. tuberculosis*

were determined during growth with different nitrogen sources, including ammonium chloride and amino acids. With asparagine as the nitrogen source, wild-type *M. tuberculosis* grew with a doubling time of 24.7 ± 0.4 h and 25.0 ± 0.4 (means \pm standard deviations) for RVW7 (Fig. 7A). Growth of both strains on ammonium chloride was slower, but with no significant difference between strains (WT, 35.4 ± 1.0 h; RVW7, 34.5 ± 1.0 h). RVW7 grew using ammonium as the sole nitrogen source, showing Ald is not essential for the synthesis of either alanine or glycine. With glycine as the nitrogen source there was no difference in growth (WT, 35.4 ± 1.1 h; RVW7, 36.7 ± 3.7 h), indicating that Ald is not essential for the breakdown of glycine. Finally, with alanine as the nitrogen source, wild-type *M. tuberculosis* grew with a doubling time of 83.2 ± 2.7 h (Fig. 7B). RVW7 was unable to grow, but the complemented strain RVW7(pAld21) was restored to growth (doubling time of 87.5 ± 2.8 h). This indicated that Ald plays an essential role in the utilization of alanine but not of glycine.

Expression of *ald*. The expression of *ald* during aerobic growth with different nitrogen sources was analyzed using quantitative real-time PCR. Expression levels were normalized to 16S RNA, with Rv0239 included as an additional control (21). Expression of *ald* was induced by growth with alanine as the sole nitrogen

TABLE 3 Apparent K_m values for His-tagged alanine dehydrogenase on different substrates

Substrate	K_m (mM)
Pyruvate	2.8
Glyoxylate	5.5
Ammonium	2.3
NADH	0.0384
Alanine	4.3
NAD ⁺	0.1328

TABLE 4 Glyoxylate and pyruvate reductive aminase activities^a

Strain, growth condition, and fraction analyzed	Sp act ($\mu\text{mol NADH oxidized/min}/\mu\text{g of protein}$) (mean \pm SD)	
	GxRA	PvRA
H37Rv		
AG, cell lysate	0.023 ± 0.004	0.010 ± 0.002
AG, culture filtrate	0.003 ± 0.001	0 ± 0.004
NRP, cell lysate	0.110 ± 0.003	0.069 ± 0.002
NRP, culture filtrate	0.001 ± 0	0 ± 0.001
Erdman		
AG, cell lysate	0.046 ± 0.001	0.014 ± 0.000
AG, culture filtrate	0.003 ± 0.003	0 ± 0.000
NRP, cell lysate	0.038 ± 0.005	0.054 ± 0.002
NRP, culture filtrate	0.003 ± 0.001	0.001 ± 0.002

^a Glyoxylate and pyruvate reductive aminase activities for H37Rv and Erdman were determined. Each sample was incubated aerobically or anaerobically, and cells and medium were then separated and analyzed for activity.




FIG 6 Subcellular location of Ald. Cell extracts and culture filtrates from aerobically growing or NRP-2 cultures were probed with an anti-Ald antibody. (A) AG, aerobically growing cells; NRP, nonreplicating persistent stage 2 cells; F, growth medium filtrate; E, cell extracts. (B) W, cell wall fraction; S, soluble fraction; M, cell membrane.

source but not by the other nitrogen sources (Fig. 8). Expression of the control gene, Rv0239, was relatively unchanged.

Enzyme activities were measured in extracts from cells grown with different nitrogen sources (Table 5). All three enzyme activities were low during growth on ammonium chloride, asparagine, or glycine. There was an approximately 5-fold induction of enzyme activity during growth with alanine.

DISCUSSION

Dehydrogenation of alanine and glycine. By using both biochemical and genetic approaches, we have shown that *ald* of *M. tuberculosis* encodes an enzyme that has both pyruvate and glyoxylate aminase activities. Inactivation of *ald* in *M. tuberculosis* resulted in loss of both activities, which were restored by the insertion of a cloned copy of *ald* into the genome. In addition, a histidine-tagged form of the protein purified from *E. coli* catalyzed both reactions.

The *gcvB* gene, annotated as a glycine dehydrogenase in the *M. tuberculosis* genome, probably encodes the P protein of the glycine cleavage system. This is based on similarity with known P proteins and because *gcvB* is in an operon with another gene, *gcvH*, that is involved in glycine catabolism. The P protein decarboxylates glycine and does not have the activity attributed to the glycine dehydrogenase. The glycine cleavage system may be responsible for utilization of glycine as a nitrogen source (Fig. 7).

Alanine dehydrogenase has been characterized from many dif-




FIG 8 Quantitative real-time PCR of *ald* grown with different nitrogen sources. RNA levels are expressed relative to stable 16S rRNA. (Insert) Expression of Rv0239. The standard deviations are shown.

ferent bacteria, but not all these enzymes have GxRA activity. This may be due to the large sequence diversity seen among these enzymes (51). There may be two classes of alanine dehydrogenases, those that can also use glyoxylate and those that cannot. Purified alanine dehydrogenases from *Anabaena cylindrica* (43) and *Streptomyces fradiae* (54) lack activity with glyoxylate. The alanine dehydrogenase of *Pseudomonas aeruginosa* has been proposed to be a glycine dehydrogenase with GxRA activity (20). Glycine produced from the enzyme was used to synthesize cyanide under hypoxic conditions (7). An *ald* knockout mutant of *P. aeruginosa* had reduced virulence in rats, showing that the enzyme, although not essential, does play a role in pathogenesis (5). In *M. tuberculosis* no cyanide was detected in growing or NRP cultures (data not shown).

Ald of *M. tuberculosis* has a greater affinity for pyruvate than for glyoxylate, but there is evidence that the GxRA reaction occurs *in vivo*. GxRA was detected in whole cells of *M. smegmatis* by isotopomer analysis (49). This activity was upregulated in response to hypoxia, while the pyruvate reductive aminase reaction was not. Each activity may be differentially regulated. In this model, the conversion of alanine to pyruvate for the purpose of nitrogen assimilation would be by alanine dehydrogenase during aerobic




FIG 7 Growth of RVW7. (A) Growth of *M. tuberculosis* in LIM with different nitrogen sources. Circles, asparagine; squares, NH_4Cl ; diamonds, glycine. WT samples, solid lines with filled symbols; RVW7, dotted lines with empty symbols. (B) Growth in LIM with alanine. Circles, WT; squares, RVW7; triangles, RVW7(pAld-Gen21). The standard deviations are shown.

TABLE 5 Enzyme activities after growth on different nitrogen sources^a

Nitrogen source	Sp act ($\mu\text{mol/min}/\mu\text{g}$ of protein) (mean \pm SD)		
	PvRA	ALD	GxRA
Asparagine	0.018 \pm 0.003	0.015 \pm 0.001	0.095 \pm 0.006
Ammonium	0.032 \pm 0.002	0.023 \pm 0.001	0.127 \pm 0.013
Glycine	0.032 \pm 0.004	0.024 \pm 0.001	0.117 \pm 0.009
Alanine	0.167 \pm 0.003	0.131 \pm 0.006	0.637 \pm 0.026

^a Enzyme activities were determined in cell extracts after growth with the indicated nitrogen source.

growth. During hypoxic dormancy, the GxRA reaction would recycle NADH for redox balancing.

Role of alanine dehydrogenase. A great deal of research has been done to characterize alanine dehydrogenases in many bacterial species, among them *M. tuberculosis*, for which a 3-dimensional structure has been constructed (1, 51). Many roles have been proposed for this enzyme, in part because of uncertainty of the main direction of the reaction. Alanine dehydrogenase may fulfill several roles, depending on the conditions and perhaps even the species.

Pyruvate reductive amination was proposed to be the main role *in vivo* serving to produce alanine (9, 24, 51) for the synthesis of peptidoglycan (2). However, RVW7 grew with NH_4Cl as the nitrogen source, indicating that in *M. tuberculosis* Ald is not essential for the synthesis of alanine or glycine.

The PvRA reaction of the alanine/glycine dehydrogenase could also be used to incorporate and store ammonia. *M. tuberculosis* assimilates ammonia by glutamine synthetase and glutamate synthetase. This is a high-affinity system that requires ATP. Glutamate dehydrogenase provides a lower-affinity non-ATP-requiring system, but this activity was not detected in *M. tuberculosis* (50). Under certain conditions, alanine dehydrogenase may fulfill an ammonium assimilation role, as has been proposed for other bacteria (6, 46). This could be important if glutamine synthetase is inhibited by high levels of alanine or glycine (9). For *Anabaena*, Ald has been proposed to synthesize alanine to store and transport nitrogen from heterocysts to vegetative cells (37).

The reverse reaction with ALD, involving the oxidative deamination of alanine to pyruvate, clearly has a role during growth with alanine as the nitrogen source in *M. tuberculosis* (Fig. 7), *M. bovis* BCG (9), and *M. smegmatis* (14). In *M. tuberculosis*, alanine dehydrogenase is required for the utilization of alanine, but not glycine, as a nitrogen source. The release of ammonium from the ALD reaction was also suggested to play a role in blocking phagosome acidification (39).

Enzyme activity and RNA levels were induced by growth with alanine as the nitrogen source (Table 5 and Fig. 8). Ammonium suppresses Ald protein levels in *M. bovis* BCG (9), while in *M. smegmatis*, ald transcription and enzyme activity were upregulated in rich media by the presence of alanine (14). Expression of alanine dehydrogenase was also upregulated by hypoxia (Table 4), which appears to be independent of the DosR/DevR regulator (56). The regulators in both cases are unknown, but a *regX3* mutant showed increased expression of *ald* during aerobic growth and reduced expression under hypoxic conditions, suggesting a role for the two-component *senX3-regX3* system (36, 40).

A role in recycling of NADH during the interruption of aerobic respiration has been proposed for Ald of *M. tuberculosis* (59, 60).

Both the GxRA and PvRA activities were induced by hypoxia (Fig. 5). A similar role was proposed for alanine dehydrogenase in *M. smegmatis* (23), and indeed, an *ald* knockout mutant was unable to grow anaerobically (14). The slower growth of an alanine dehydrogenase mutant in *Anabaena* could also be due to impairment of NADH recycling (37).

Additional roles for alanine dehydrogenase are possible. These include the utilization and detoxification of propionate or glyoxylate. The main energy source for *M. tuberculosis in vivo* is thought to be fatty acids. β -Oxidation of fatty acids containing an even number of carbons produces acetyl coenzyme A (CoA), while β -oxidation of fatty acids with odd numbers of carbons and cholesterol produces both acetyl-CoA and propionyl-CoA. Propionyl-CoA is converted to methylisocitrate in the methylcitrate cycle and then to pyruvate by isocitrate lyase (18). Acetyl-CoA enters the glyoxylate cycle and is converted by isocitrate lyase to glyoxylate. Both of these products, pyruvate and glyoxylate, are substrates for alanine dehydrogenase. This enzyme may connect these two cycles, regulating glyoxylate and propionate levels.

Location. Alanine dehydrogenase was detected only in association with the cells of *M. tuberculosis*, not in the extracellular medium. Specifically, most of the protein was isolated with the membrane and a lesser amount in the cytosol. Neither alanine dehydrogenase protein (Fig. 6) nor enzymatic activity (Table 4) was detected in the culture filtrate. In contrast, others have detected protein (2, 28, 34, 42) and activity (39) in the extracellular medium of growing cultures of *M. tuberculosis*. In *Mycobacterium marinum* and in *M. bovis* expressing the *M. tuberculosis ald*, alanine dehydrogenase was secreted (9). We used DTA medium while the other reports utilized Sautons medium for growth, but the difference is probably not due to the medium, as a comprehensive analysis of proteins exported from H37Rv also cultured in Sautons medium did not detect alanine dehydrogenase (31).

GxRA and PvRA activities were detected in cell filtrates, but not culture filtrates, of both the H37Rv and Erdman strains. Proteome analysis had detected this enzyme only in H37Rv and not the Erdman strain (25). The Erdman strain expressed higher levels during aerobic growth (Table 4).

Malate synthase, which is involved in the glyoxylate shunt, and the glycolytic enzyme fructose-1,6-bisphosphate aldolase are exported to the cell surface of *M. tuberculosis* (10, 26). Both of these enzymes and alanine dehydrogenase lack signal sequences. In addition to their metabolic function, they are involved in host-pathogen interactions. Alanine dehydrogenase may also play such a role, which would explain its membrane location. Alternatively, GxRA and ALD activities may be regulated by an interaction with membrane components of the respiratory chain.

Importance of Ald in pathogenesis. It is intriguing that alanine dehydrogenase is important in other bacteria that have specialized persistence programs, such as sporulation. In *Myxococcus xanthus*, an *ald* mutant showed delayed aggregation and reduced levels of sporulation (57). An *ald* mutant of *Bacillus subtilis* was defective for sporulation (47). This was partially complemented by the addition of pyruvate, suggesting that the role of alanine dehydrogenase is to provide pyruvate as an energy source.

The *ald* gene of *M. bovis* contains a single nucleotide deletion and therefore lacks this enzyme (9). Also absent due to a point mutation is pyruvate kinase, suggesting there are major differences in central metabolism between *M. bovis* and *M. tuberculosis* (8, 55). Despite these changes, *M. bovis* can produce disease in

humans. *M. bovis* BCG expressing the *M. tuberculosis* *ald* showed similar survival in both macrophages and mice (44). Alanine dehydrogenase is probably not essential for pathogenesis in *M. tuberculosis* but may still play an important role.

ACKNOWLEDGMENTS

We thank Sylors Chem and Karen Kelley for valuable discussions and review of the manuscript, Sandy Sudberg for technical help, and Sheldon Broedel at AthenaES for protein purification.

The following reagent was obtained through the NIH Biodefense and Emerging Infections Research Resources Repository, NIAID, NIH: monoclonal anti-*Mycobacterium tuberculosis* Ald (gene Rv2780), clone IT-46 (HBT10) (produced *in vitro*) (NR-13651). This study was supported by the Medical Research Services of the U.S. Department of Veterans Affairs.

REFERENCES

- Agren D, et al. 2008. Three-dimensional structures of apo- and holo-L-alanine dehydrogenase from *Mycobacterium tuberculosis* reveal conformational changes upon coenzyme binding. *J. Mol. Biol.* 377:1161–1173.
- Andersen AB, Andersen P, Ljungqvist L. 1992. Structure and function of a 40,000-molecular-weight protein antigen of *Mycobacterium tuberculosis*. *Infect. Immun.* 60:2317–2323.
- Awasthy D, et al. 2009. Inactivation of the *ilvB1* gene in *Mycobacterium tuberculosis* leads to branched-chain amino acid auxotrophy and attenuation of virulence in mice. *Microbiology* 155:2978–2987.
- Beste DJV, et al. 2011. ¹³C metabolic flux analysis identifies an unusual route for pyruvate dissimilation in mycobacteria which requires isocitrate lyase and carbon dioxide fixation. *PLoS Pathog.* 7:1–18.
- Boulette ML, et al. 2009. Characterization of alanine catabolism in *Pseudomonas aeruginosa* and its importance for proliferation *in vivo*. *J. Bacteriol.* 191:6329–6334.
- Caballero FJ, Cárdenas J, Castillo F. 1989. Purification and properties of L-alanine dehydrogenase of the phototrophic bacterium *Rhodobacter capsulatus* E1F1. *J. Bacteriol.* 171:3205–3210.
- Castric PA. 1983. Hydrogen cyanide production by *Pseudomonas aeruginosa* at reduced oxygen levels. *Can. J. Microbiol.* 29:1344–1349.
- Chavadi S, et al. 2009. Global effects of inactivation of the pyruvate kinase gene in the *Mycobacterium tuberculosis* complex. *J. Bacteriol.* 191:7545–7553.
- Chen JM, Alexander DC, Behr MA, Liu J. 2003. *Mycobacterium bovis* BCG vaccines exhibit defects in alanine and serine catabolism. *Infect. Immun.* 71:708–716.
- de la Paz Santangelo M, et al. 2011. Glycolytic and non-glycolytic functions of the fructose-1,6-bisphosphate aldolase of *Mycobacterium tuberculosis*, an essential enzyme produced by replicating and non-replicating Bacilli. *J. Biol. Chem.* 286:40219–40231.
- Desjardin LE, Hayes LG, Sohaskey CD, Wayne LG, Eisenach KD. 2001. Microaerophilic induction of the alpha-crystallin chaperone protein homologues (*hspX*) mRNA of *Mycobacterium tuberculosis*. *J. Bacteriol.* 183:5311–5316.
- Emsley P, Lohkamp B, Scott WG, Cowtan K. 2010. Features and development of *Coot*. *Acta Crystallogr. D Biol. Crystallogr.* 66:486–501.
- Eng JK, McCormack AL, Yates JR, III. 1994. An approach to correlate tandem mass spectral data of peptides with amino acid sequences in a protein database. *J. Am. Soc. Mass Spec.* 5:976–989.
- Feng Z, Cáceres NE, Sarath G, Barletta RG. 2002. *Mycobacterium smegmatis* L-alanine dehydrogenase (Ald) is required for proficient utilization of alanine as a sole nitrogen source and sustained anaerobic growth. *J. Bacteriol.* 184:5001–5010.
- Goldman DS. 1959. Enzyme systems in the mycobacteria. *Bioch. Biophys. Acta* 34:527–539.
- Goldman DS, Wagner MJ. 1962. Enzyme systems in the mycobacteria XIII glycine dehydrogenase and the glyoxylic acid cycle. *Biochim. Biophys. Acta* 65:297–306.
- Gordhan BG, et al. 2002. Construction and phenotypic characterization of an auxotrophic mutant of *Mycobacterium tuberculosis* defective in L-arginine biosynthesis. *Infect. Immun.* 70:3080–3084.
- Gould TA, van de Langemheen H, McKinney JD, Sacchettini JC. 2006. Dual role of isocitrate lyase 1 in the glyoxylate and methylcitrate cycles in *Mycobacterium tuberculosis*. *Mol. Microbiol.* 61:940–947.
- Gu S, et al. 2003. Comprehensive proteomic profiling of the membrane constituents of a *Mycobacterium tuberculosis* strain. *Mol. Cell. Proteomics* 2:1284–1296.
- Hagins JM, Locy R, Silo-Suh L. 2009. Isocitrate lyase supplies precursors for hydrogen cyanide production in a cystic fibrosis isolate of *Pseudomonas aeruginosa*. *J. Bacteriol.* 191:6335–6339.
- Honaker RW, Leistikow RL, Bartek IL, Voskuil MI. 2009. Unique roles of DosT and DosS in DosR regulon induction and *Mycobacterium tuberculosis* dormancy. *Infect. Immun.* 77:3258–3263.
- Hondalus MK, et al. 2000. Attenuation of and protection induced by a leucine auxotroph of *Mycobacterium tuberculosis*. *Infect. Immun.* 68:2888–2898.
- Hutter B, Dick T. 1998. Increased alanine dehydrogenase activity during dormancy in *Mycobacterium smegmatis*. *FEMS Microbiol. Lett.* 167:7–11.
- Hutter B, Singh M. 1999. Properties of the 40 kDa antigen of a *Mycobacterium tuberculosis*, a functional L-alanine dehydrogenase. *Biochem. J.* 343:669–672.
- Jungblut PR, et al. 1999. Comparative proteome analysis of *Mycobacterium tuberculosis* and *Mycobacterium bovis* BCG strains: towards functional genomics of microbial pathogens. *Mol. Microbiol.* 33:1103–1117.
- Kinhikar AG, et al. 2006. *Mycobacterium tuberculosis* malate synthase is a laminin-binding adhesin. *Mol. Microbiol.* 60:999–1013.
- Lim A, Eleuterio M, Hutter B, Murugasu-Oei B, Dick T. 1999. Oxygen depletion induced dormancy in *Mycobacterium bovis* BCG. *J. Bacteriol.* 181:2252–2256.
- Ljungqvist L, Worsaae A, Heron I. 1988. Antibody responses against *Mycobacterium tuberculosis* in 11 strains of inbred mice: novel monoclonal antibody specificities generated by fusions, using spleens from BALB. B10 and CBA/J mice. *Infect. Immun.* 56:1994–1998.
- Lyon RH, Hall WH, Costas-Martinez C. 1970. Utilization of amino acids during growth of *Mycobacterium tuberculosis* in rotary cultures. *Infect. Immun.* 1:513–520.
- Mahenthiralingam E, et al. 1998. Site-directed mutagenesis of the 19-kilodalton lipoprotein antigen reveals no essential role for the protein in the growth and virulence of *Mycobacterium intracellulare*. *Infect. Immun.* 66:3626–3634.
- Malen H, Berven FS, Fladmark KE, Wiker HG. 2007. Comprehensive analysis of exported proteins from *Mycobacterium tuberculosis* H37Rv. *Proteomics* 7:1702–1718.
- McAdam RA, et al. 1995. *In vivo* growth characteristics of leucine and methionine auxotrophic mutants of *Mycobacterium bovis* BCG generated by transposon mutagenesis. *Infect. Immun.* 63:1004–1012.
- McDonough JA, et al. 2008. Identification of functional Tat signal sequences in *Mycobacterium tuberculosis* proteins. *J. Bacteriol.* 190:6428–6438.
- Ortalo-Magné A, et al. 1995. Molecular composition of the outermost capsular material of the tubercle bacillus. *Microbiology* 141:1609–1620.
- Parish T. 2003. Starvation survival response of *Mycobacterium tuberculosis*. *J. Bacteriol.* 185:6702–6706.
- Parish T, Smith DA, Roberts G, Betts J, Stoker NG. 2003. The *senX3-regX3* two-component regulatory system of *Mycobacterium tuberculosis* is required for virulence. *Microbiology* 149:1423–1435.
- Pernil R, Herrero A, Flores E. 2010. Catabolic function of compartmentalized alanine dehydrogenase in heterocyst-forming cyanobacterium *Anabaena* sp. strain PCC 7120. *J. Bacteriol.* 192:5165–5172.
- Quandt J, Hynes MF. 1993. Versatile suicide vectors which allow direct selection for gene replacement in Gram-negative bacteria. *Gene* 127:15–21.
- Raynaud C, Etienne G, Peyron P, Lanéelle M-A, Daffé M. 1998. Extracellular enzyme activities potentially involved in the pathogenicity of *Mycobacterium tuberculosis*. *Microbiology* 144:577–587.
- Roberts G, Vadrevu I, Madiraju MV, Parish T. 2011. Control of CydB and GltA1 expression by the SenX3 RegX3 two component regulatory system of *Mycobacterium tuberculosis*. *PLoS One* 6:1–9.
- Rosenkrands I, et al. 2002. Hypoxic response of *Mycobacterium tuberculosis* studied by metabolic labeling and proteome analysis of cellular and extracellular proteins. *J. Bacteriol.* 184:3485–3491.
- Rosenkrands I, et al. 2000. Mapping and identification of *Mycobacterium tuberculosis* proteins by two-dimensional gel electrophoresis, microsequencing and immunodetection. *Electrophoresis* 21:948.
- Rowell P, Stewart DP. 1976. Alanine dehydrogenase of the N₂-fixing blue green alga, *Anabaena cylindrica*. *Arch. Microbiol.* 107:115–124.
- Scandurra GM, Ryan AA, Pinto R, Britton WJ, Triccas JA. 2006.

- Contribution of L-alanine dehydrogenase to *in vivo* persistence and protective efficacy of the BCG vaccine. *Microbiol. Immunol.* **50**:805–810.
45. Schaefer WB, Marshak A, Burkhart B. 1949. The growth of *Mycobacterium tuberculosis* as a function of its nutrients. *J. Bacteriol.* **58**:549–563.
 46. Schuffenhauer G, Schröder T, Andreesen JR. 1999. Morpholine-induced formation of L-alanine dehydrogenase activity in *Mycobacterium* strain HE5. *Arch. Microbiol.* **171**:417–423.
 47. Siranosian KJ, Ireton K, Grossman AD. 1993. Alanine dehydrogenase (*ald*) is required for normal sporulation in *Bacillus subtilis*. *J. Bacteriol.* **175**:6789–6796.
 48. Sohaskey CD, Wayne LG. 2003. Role of *narK2X* and *narGHJ1* in hypoxic upregulation of nitrate reduction by *Mycobacterium tuberculosis*. *J. Bacteriol.* **185**:7247–7256.
 49. Tang YJ, et al. 2009. Central metabolism in *Mycobacterium smegmatis* during the transition from O₂-rich to O₂-poor conditions as studied by isotopomer-assisted metabolite analysis. *Biotechnol. Lett.* **31**:1233–1240.
 50. Tian J, Bryk R, Itoh M, Suematsu M, Nathan C. 2005. Variant tricarboxylic acid cycle in *Mycobacterium tuberculosis*: identification of α -ketoglutarate decarboxylase. *Proc. Natl. Acad. Sci. U. S. A.* **102**:10670–10675.
 51. Tripathi SM, Ramachandran R. 2008. Crystal structures of the *Mycobacterium tuberculosis* secretory antigen alanine dehydrogenase (Rv2780) in apo and ternary complex forms captures “open” and “closed” enzyme conformations. *Proteins* **72**:1089–1095.
 52. Tullius MV, Harth G, Horwitz MA. 2003. Glutamine synthetase GlnA1 is essential for growth of *Mycobacterium tuberculosis* in human THP-1 macrophages and guinea pigs. *Infect. Immun.* **71**:3927–3936.
 53. Usha V, Jayaraman R, Toro JC, Hoffner SE, Das KS. 2002. Glycine and alanine dehydrogenase activities are catalyzed by the same protein in *Mycobacterium smegmatis*: upregulation of both activities under microaerophilic adaptation. *Can. J. Microbiol.* **48**:7–13.
 54. Vancura A, et al. 1989. Alanine dehydrogenase from *Streptomyces fradiae*. *Eur. J. Biochem.* **179**:221–227.
 55. van Keulen G, Alderson J, White J, Sawers RG. 2005. Nitrate respiration in the actinomycete *Streptomyces coelicolor*. *Biochem. Soc. Trans.* **33**:210–212.
 56. Voskuil MI, et al. 2002. Inhibition of respiration by nitric oxide induces a *Mycobacterium tuberculosis* persistence program. *J. Exp. Med.* **198**:705–713.
 57. Ward MJ, Lew H, Zusman DR. 2000. Disruption of *aldA* influences the developmental process in *Myxococcus xanthus*. *J. Bacteriol.* **182**:546–550.
 58. Wayne LG, Hayes LG. 1996. An in vitro model for sequential study of shiftdown of *Mycobacterium tuberculosis* through two stages of nonreplicating persistence. *Infect. Immun.* **64**:2062–2069.
 59. Wayne LG, Lin K-Y. 1982. Glyoxylate metabolism and adaptation of *Mycobacterium tuberculosis* to survival under anaerobic conditions. *Infect. Immun.* **37**:1042–1049.
 60. Wayne LG, Sohaskey CD. 2001. Nonreplicating Persistence of *Mycobacterium tuberculosis*. *Annu. Rev. Microbiol.* **55**:139–163.