

HOXB13 Homeodomain Protein Suppresses the Growth of Prostate Cancer Cells by the Negative Regulation of T-Cell Factor 4

Chaeyong Jung, Ran-Sook Kim, Sang-Jin Lee, et al.

Cancer Res 2004;64:3046-3051.

Updated version Access the most recent version of this article at:
<http://cancerres.aacrjournals.org/content/64/9/3046>

Cited Articles This article cites by 37 articles, 14 of which you can access for free at:
<http://cancerres.aacrjournals.org/content/64/9/3046.full.html#ref-list-1>

Citing articles This article has been cited by 15 HighWire-hosted articles. Access the articles at:
<http://cancerres.aacrjournals.org/content/64/9/3046.full.html#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, contact the AACR Publications Department at permissions@aacr.org.

HOXB13 Homeodomain Protein Suppresses the Growth of Prostate Cancer Cells by the Negative Regulation of T-Cell Factor 4

Chaeyong Jung,¹ Ran-Sook Kim,¹ Sang-Jin Lee,¹ Chihuei Wang,⁴ and Meei-Huey Jeng^{2,3}

Departments of ¹Urology, ²Microbiology and Immunology, and ³Medicine, Walther Oncology Center, Indiana University, Indianapolis, Indiana, and ⁴the Orthopedic Research Center and Department of Biology, Kaohsiung Medical University, Taiwan, Republic of China

ABSTRACT

In prostate gland, HOXB13 is highly expressed from the embryonic stages to adulthood. However, the function of HOXB13 in normal cell growth and tumorigenesis is not yet known. We investigated the role of HOXB13 and mechanism by which it functions in HOXB13-negative cells. Expression of HOXB13 was forced in HOXB13-negative PC3 prostate cancer cells using a liposome-mediated gene transfer approach. Compared with the control clones, HOXB13-expressing PC3 cells exhibited significant inhibition of *in vitro* and *in vivo* cell growth with G₁ cell cycle arrest mediated by the suppression of cyclin D1 expression. Because cyclin D1 is mainly regulated by β -catenin/T-cell factor (TCF), TCF-4 response element was used in a reporter gene transcription assay, demonstrating that HOXB13 significantly inhibits TCF-4-mediated transcriptional activity in both prostate and nonprostate cells. This inhibition occurred in a dose-responsive manner and was specific to TCF-4 response element. Western blot analysis demonstrated that HOXB13 down-regulates the expression of TCF-4 and its responsive genes, *c-myc* and *cyclin D1*. HOXB13 also suppressed the activity of natural *c-myc* promoter. This study suggests that HOXB13, a transcription factor, functions as a cell growth suppressor by negatively regulating the expression of TCF-4, which eventually provides negative signals for cell proliferation. This observation will provide valuable insight into the molecular basis of prostate tumorigenesis.

INTRODUCTION

Prostate cancer (PCa) is the second leading cancer killer in North American males. The mechanisms controlling proliferation, differentiation, and tumorigenesis in the prostate are largely unknown, and therapeutic efforts to conquer this malignant disease have been limited. There have been attempts to identify genes that might be involved in prostatic development and carcinogenesis. Many studies have concentrated on the activation of oncogenes and inactivation of tumor suppressor genes. Several homeobox genes have been also described in the normal and malignant growth of prostate epithelial cells. *Nkx3.1* is the best-studied homeobox gene in PCa. Expression of *Nkx3.1* is very important for prostatic organogenesis and maintaining the normal growth of epithelial cells (1). *Nkx3.1* knockout mice display reductions in prostatic duct tips and disrupted secretory function. At the same time, *Nkx3.1* deletion mutants display a high incidence of prostatic intraepithelial neoplasia (2, 3). The loss of *NKX3.1* is more common in tumors than in benign prostatic hyperplasia. Almost 80% of metastatic tumors lose *NKX3.1* expression (3, 4).

HOX genes are known to regulate axial regional specification during embryonic development. They also have developmental roles in male accessory sexual organs, including the prostate (5–7). The *Hox-13* paralogue is especially important to prostate development. All

Hox-13 genes except *Hoxc13* are expressed in the prostate during embryonic development, but *Hoxa13* and *Hoxd13* are no longer expressed after the embryonic stage (5, 8). *Hoxa13* and/or *Hoxd13*-deficient mice showed morphological abnormalities in male accessory sex organs, including the seminal vesicle and prostate (5, 6, 9). *Hoxb13*, the last identified vertebrate *Hox* gene, has limited expression in the caudal extent of the spinal cord, tail bud, and urogenital sinus (10). In the prostate gland, embryologically derived from the urogenital sinus, expression of *Hoxb13* is not regulated by androgen, in contrast to *Nkx3.1* (11). Recently, *Hoxb13* loss-of-function mutations showed overgrowth in all major structures derived from the tail bud (12) and malformation of the ventral prostate ducts, including complete loss of secretory proteins (13). The phenotype demonstrated in *Hoxb13* mutant mice was similar to *Nkx3.1* mutant mice, the development of swollen prostates in older mutant mice.

Although *Hoxb13* expression is maintained throughout adulthood, only limited information is available on the biological function of HOXB13 in normal prostate development and human prostate tumorigenesis. We characterized the biological function of HOXB13 using the PC3 PCa cell line, which does not express this gene.

MATERIALS AND METHODS

Cell Lines and Tissues. Human PCa cell lines PC3 and CWR22RV were cultured in RPMI media (Invitrogen) supplemented with 5% FCS at 37°C in an atmosphere containing 5% CO₂ as described previously (14). MDA PCa 2b PCa cells were grown in Biological Research Faculty and Facility (BRFF)-HPC1 medium (AthenaES, Baltimore, MD) with 20% fetal bovine serum (FBS). CV-1 monkey kidney and 293 human embryonic renal cells were maintained in DMEM containing 10% FBS. All cultures were fed with fresh medium every 3–4 days.

Construction of HOXB13 Expression Vectors. *HOXB13* full-length cDNA was obtained by reverse transcription-PCR (RT-PCR) using human prostate total RNA purchased from Clontech. Amplified *HOXB13* cDNA was first cloned into pFLAG-cytomegalovirus (CMV)2 vector (Sigma) to be linked to an NH₂-terminal FLAG tag. The resulting pFLAG-*HOXB13* vector was confirmed by sequencing analysis and transient transfection into 293 cells, followed by Western blot analysis. FLAG-*HOXB13* (*fHOXB13*) was then excised from pFLAG-*HOXB13* vector and subcloned into multiple cloning sites of CMV-driven pIRES-neo expression vector (Clontech). An adenoviral *HOXB13* expression construct was also developed by modifying adenoviral vector pAd1020sfidA (OD260, Inc.). Briefly, the expression cassette CMV-multiple cloning sites-pA-CMV-GFP-pA was excised from pAdTrack-CMV (a generous gift from Dr. Bert Vogelstein) and ligated into pAd1020sfidA vector. Then, *fHOXB13* was cloned into the multiple cloning sites of modified pAd1020sfidACMVGFP, generating pAd1020GFP-*fHOXB13*. The left arm of the adenovirus with the expression cassette was cut out by digestion with *Sfi*I and *Pac*I and ligation into AdenoZapsfi.2 (OD260, Inc.) in the presence of *Pac*I. Purified DNA was transfected to 293 cells using LipofectAMINE 2000 (Invitrogen), and virus was produced and purified as described previously (15).

Transfection and Colony Formation. To determine the effects of HOXB13 on PCa cell growth, PC3 (1×10^5) cells were plated in P60 dishes and transfected with either pIRES-neo as a control vehicle or pIRES-*fHOXB13* vector using *N*-(1-(2,3-dioleoyloxy)propyl)-*N,N,N*-trimethylammoniummethyl sulfate transfection reagent (Roche Molecular Biochemicals) as described by the manufacturer. Cells were selected in the presence of geneticin (200 μ g/ml final concentration; Cellgro). When visible colonies (~200 cells) were formed, the number of colonies was counted. For the generation of stable transfectants, PC3

Received 8/21/03; revised 2/24/04; accepted 2/27/04.

Grant support: Grants from the Indiana University Biomedical Research Fund 22-88-430, Thomson Prostate Cancer Research Fund, and Albert James Hackl Urologic Oncology Fellowship.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Requests for reprints: Chaeyong Jung, Department of Urology, Indiana University, 1001 West 10th Street, OPW320, Indianapolis, IN 46202. Phone: (317) 278-7617; Fax: (317) 278-3432; E-mail: chjung@iupui.edu.

cells were seeded at 1×10^6 cells onto a P100 tissue culture dish and transfected as described above. The resulting neomycin-resistant clones were ring cloned and expanded.

Western Blot. Cells were grown $\leq 80\%$ confluency in P60 culture dishes containing 5% FBS-T media. Cells were then lysed in protein extraction buffer ($1 \times$ TBS, 1% NP40, 0.5% sodium deoxycholate, 0.1% SDS, and protease inhibitors). Total cell lysates (20 μ g) were loaded onto 10% Bis-Tris gel (Invitrogen) and separated using a Novex electrophoresis system (Invitrogen). After proteins were transferred to polyvinylidene difluoride membrane, the expression of FLAG-tagged HOXB13 was detected using FLAG M2 monoclonal antibody (Sigma) followed by the enhanced chemiluminescence detection system (Pierce).

RT-PCR. Total RNA from each cell line was extracted using the Ultraspec RNA isolation system (Biotecx Laboratories, Inc., Houston, TX) as described previously (16). cDNA was obtained from 1 μ g of corresponding total RNA using Moloney murine leukemia virus reverse transcriptase. *HOXB13* primers (5'-ccccactgagttgctctatc-3' and 5'-gcctctgtctctgtgatgaac-3') are designed to specifically amplify a coding region of 369–739. Cyclin D1 (5'-tgtgctgcgaagtggaaacc-3' and 5'-aaatcgtgcggggtcattgc-3') and T-cell factor 4 (TCF-4; 5'-tcccaccatcatagctacac-3' and 5'-tcgtgtctctctctggacag-3') primers were also constructed. Primers for β -actin were used to show equal loading (5'-gcaccacacctctacatgagc-3' and 5'-tagcacagctgtgatgacagc-3'). To be close to the linear range for PCR, equal amounts of RNA and cDNA products were applied with a minimum number of amplification cycles (≤ 25 cycles).

3-(4,5-Dimethylthiazol-2-yl)-2,5-Diphenyltetrazolium Bromide Cell Proliferation Assay. Stably transfected HOXB13-PC3 cells were plated at 1.5×10^3 cells/well into 96-well tissue culture plates and incubated at 37°C in a humidified atmosphere containing 5% CO₂. At the indicated time points, 20 μ l of 5 mg/ml 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide solution were added to the cell cultures and incubated for 2 h at 37°C. The reaction was stopped by adding 100 μ l of extraction buffer [50% formamide and 10% SDS (pH 4.7)]. After overnight incubation at 37°C, the absorbance at 570 nm was measured using a Spectra Microplate Reader (Molecular Devices).

Flow Cytometric Analysis. Cells were grown in 5% FBS-T media containing 400 μ g/ml geneticin for 72 h, harvested, fixed, and treated with RNase. Cells were then stained with propidium iodide (50 μ g/ml) and analyzed using a FACScan Flow Cytometer and the Cell Quest program.

Tumor Growth in Vivo. Male 8-week-old athymic nude mice (BALB/c-nu/nu) were obtained from Harlan. Mice were kept in groups of three according to the institutional guidelines for animal welfare. To characterize the tumorigenicity of *HOXB13*-transfected PC3 clones *in vivo*, HOXB13-PC3 clones 1 and 5 were injected into nude mice s.c. A vector-only transfected clone and wild-type PC3 cells were used as controls. Three animals were used for each clone. Cells (1×10^6 in 0.1 ml of T-media containing 5% FBS) were injected s.c. into the back of each nude mouse. When the tumor was palpable, it was measured using a caliper each week. Tumor volumes were calculated by the formula $TV \text{ (mm}^3\text{)} = (L \times W^2)/2$, where L is the longest dimension of the tumor (in millimeters), and W is the shortest dimension of the tumor (in millimeters).

Reporter Gene Transcription Analysis. To investigate the involvement of HOXB13 in β -catenin/TCF-mediated transcriptional activity, pTCF-4 response element (TCF-4RE)-luc and a constitutively active form of β -catenin were constructed. pTCF-4RE-luc contains four copies of the TCF-4REs derived from cyclin D1 promoter. The constitutive form of β -catenin is mutated at amino acids 41 and 45 to prevent phosphorylation. pCDNA-*hNKX3.1* and pBV-cmyc-luc were generously provided by Drs. Charles Bieberich and Bert Vogelstein, respectively. For the transient transfection assay, $\sim 1 \times 10^5$ cells were plated in a 24-well plate 16 h before transfection. Transfections were carried out using LipofectAMINE 2000 (Invitrogen) for all of the cells as described by the manufacturer's protocol. Six h after transfection, the cells were washed and fed with medium containing 5% FBS. Cells were incubated for 36 h, and luciferase activity was measured as relative light units normalized by β -galactosidase activity. Each transfection experiment was done in triplicate.

Adenoviral Infections. For the HOXB13-induced cell killing assay, 1×10^5 cells were seeded onto six-well plate culture dishes. The next day, cells were infected with either Ad-green fluorescent protein (GFP) control virus or Ad-GFP-HOXB13 virus at a multiplicity of infection of 100. Infectivity and cell viability were carefully monitored by fluorescence microscopy

for 10 days. For Western blot, cells were grown to 70% confluence and infected with recombinant adenovirus (100 multiplicity of infection). Thirty-six h after the infection, cells were collected to isolate the cell lysates, as described above.

Statistical Analysis. Where appropriate, the data are shown as means \pm SE. To access the statistical significance of observed differences, we used the Wald test. All statistical tests were two sided, and P s < 0.01 were considered to be statistically significant.

RESULTS

To investigate potential growth-regulating functions of HOXB13 in PC3 PCa cells, cells were transfected with either pIRES-neo (Clontech) or pIRES-*fHOXB13* constitutive expression vector, and transfected cells were selected by geneticin (Cellgro). After 3 weeks of selection, isolated colonies (~ 200 –300 cells/colony) were counted. As shown in Fig. 1A, introduction of exogenous HOXB13 in HOXB13-negative PC3 cells resulted in statistically significant growth suppression, with a 20–40-fold decrease in the number of geneticin-resistant clones compared with neo control clones (Wald test; $P < 0.0001$). Growth suppression in PC3 cells was accompanied by dramatic changes in cell morphology to a gigantic cobblestone appearance (Fig. 1B), characteristic of terminal differentiated cells. By comparison, HOXB13 expression is down-regulated in developing dermis compared with adult skin, and the loss of HOXB13 causes adult skin to adopt a less differentiated and more proliferative mode (17, 18).

Thirteen neomycin-resistant clones were expanded and analyzed for HOXB13 expression using FLAG M2 antibody (Sigma). Only five clones expressed low levels of HOXB13 protein compared with the transiently transfected 293 cells, most likely because highly HOXB13-expressed clones were not expandable because of HOXB13-induced cell toxicity (Fig. 2A). *HOXB13* mRNA levels were also verified by RT-PCR analysis, which also showed no ex-





Fig. 1. HOXB13 reduces colony formation in PC3 cells. Cells were transfected with pIRES-neo or pIRES-*fHOXB13* followed by neomycin selection. In A, surviving clones were counted. B, morphological changes in transfected cells. Forced expression of HOXB13 changed the morphology of PC3 cells.

Fig. 2. Forced expression of HOXB13 inhibits growth of PC3 cells by cell cycle arrest. In A, stable transfectants expressing fHOXB13 (clones 1–5) were analyzed by Western blot. fHOXB13 expression was not detected in nontransfected PC3 (WT) or PC3 cells transfected with vector alone (neo), but was very strong in 293 cells transiently transfected with pFLAG-HOXB13. Expression of *HOXB13* mRNA was measured by reverse transcription-PCR using total RNA. In B, clones of transfected cells were analyzed by 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide proliferation assay. Absorbance is shown as $\pm 95\%$ confidence interval ($n = 8$). In C, selected HOXB13-expressing clones were injected into athymic mice, and tumor volume was determined using a caliper; bars, \pm confidence interval ($n = 3$). In D, flow cytometric analysis was performed on transfected cells using propidium iodide. Values represent the percentage of cells in each phase of the cell cycle. Cells (10,000) were counted for each experiment. In E, reverse transcription-PCR analysis demonstrated that HOXB13 down-regulates the expression of cyclin D1 RNA in HOXB13-PC3 cells.



pression of *HOXB13* in parental PC3 cells using *HOXB13*-specific primers. HOXB13-PC3 clones were further analyzed by 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide *in vitro* cell proliferation assays. As shown in Fig. 2B, all HOXB13-expressing cells grew much more slowly than the neo clone (Wald test; $P < 0.0001$). Although clone 4 showed significant growth inhibition, this clone was ruled out, because it seemed to have an abnormal number of chromosomes (duplication of chromosomes is commonly seen in cancer cells). To reiterate the growth-inhibiting effect of HOXB13 on tumor growth *in vivo*, HOXB13-transfected and control cells were injected s.c. into the backs of nude mice. Compared with the control groups, including neo-transfected clone and wild-type PC3 cells, HOXB13-transfected clones showed a dramatic decrease in tumor volumes at 5 weeks after implantation (Fig. 2C; Wald test; $P = 0.0014$). At 5 weeks after implantation, the tumor volumes of HOXB13-transfected clones 1 and 5 were $92 \pm 18 \text{ mm}^3$ and $140 \pm 45 \text{ mm}^3$, respectively. On the other hand, wild-type PC3 and PC3-neo controls were $684 \pm 117 \text{ mm}^3$ and $522 \pm 171 \text{ mm}^3$, respectively.

We then investigated whether inhibition of cell growth was caused by apoptosis or cell cycle arrest. Flow cytometric analysis demonstrated that the growth-inhibiting effect was caused by significant abnormalities in cell cycle distribution. The majority of clones (three of four) showed a dramatic increase in the G₀-G₁ cell population (Fig. 2D), which may ultimately drive the cells into death. That there was no alteration of clone 3 cell distribution seems to be attributable to a low expression of HOXB13, as shown in Fig. 2A. Not all HOXB13-PC3 clones showed an alteration in apoptotic activity (data not shown). These results suggest that HOXB13 plays a role in the suppression of cell growth in highly malignant and metastatic PC3

Pca cells through cell cycle arrest in the G₁ phase. Because cyclin D1 is integral for accelerating the G₁ phase of the cell cycle (19), we used RT-PCR to test whether the expression of cyclin D1 was altered in HOXB13-expressing PC3 cells. Surprisingly, cyclin D1 expression was down-regulated in most clones (Fig. 2E). We failed to further characterize the function of HOXB13, because not all HOXB13-PC3 cells survived in culture, probably because of the cytotoxic effect of HOXB13.

We studied whether HOXB13 can inhibit cyclin D1 promoter activity. Because β -catenin/TCF-4 is an important transcription factor for cyclin D1 promoter, we used a luciferase reporter plasmid consisting of four copies of a TCF-4RE to test whether HOXB13 regulates this artificial promoter activity. Because both LNCaP and PC3 cells did not respond well to β -catenin-activated TCF signaling, as reported previously (data not shown; Ref. 20), CWR22RV and MDA PCa 2b PCa cells were used instead. CV-1 monkey kidney cells were also used for their high transfection efficiency and to simplify the reporter transcription assay. CV-1 cells did not require exogenous β -catenin to activate β -catenin/TCF signaling (data not shown). Reporter gene transcription analysis showed that HOXB13 significantly suppressed β -catenin/TCF activity in both PCa and CV-1 cells (Fig. 3A). HOXB13-mediated suppression was accomplished in a dose-responsive manner in CV-1 (Fig. 3B) but was not seen in the other promoters, including SV40, RSV, TATA, and CMV (Fig. 3C). Instead, HOXB13 moderately increased transcriptional activity in these promoters. Additionally, overexpression of another homeobox protein, NKX3.1, did not suppress the activity of β -catenin/TCF (Fig. 4D). These results suggest that HOXB13 specifically suppresses β -catenin/TCF-4-mediated transcription.




Fig. 3. HOXB13 specifically suppresses T-cell factor 4 transcriptional activity. Cells were transfected with a pTCF-4 response element reporter construct (100 ng), expression constructs [pFLAG-cytomegalovirus (CMV), pFLAG-HOXB13, pIRES- β -catenin; 100 ng] as listed, and 20 ng of pCMV- β gal. The total amount of DNA was brought to ≤ 300 ng with pFLAG-CMV. Luciferase assays were performed at 48 h post-transfection. In A, each cell was tested for the effect of HOXB13 on T-cell factor 4 response. In B, other promoters (100 ng for each) were tested for HOXB13-induced transcriptional suppression in CWR22RV cells. In C, CV-1 cells were used for the dose response activity of HOXB13. In D, *NKX3.1* was cotransfected with pTCF-4 response element reporter construct in CWR22RV cells. In E, various amounts of β -catenin were cotransfected with pFLAG-HOXB13 into CV-1 cells. Bars, \pm SE ($n = 3$).

To test whether β -catenin acts as a modulator mediating HOXB13 suppressor function in this signaling pathway, a constitutively active form of β -catenin was cotransfected. The addition of this β -catenin did not restore the HOXB13-induced inhibition of TCF activity in CV-1 cells (Fig. 3E), suggesting that suppression of β -catenin/TCF signaling is not mediated through the alteration of β -catenin expression. Interaction of HOXB13 with β -catenin was not seen (data not shown).

To further investigate the mechanistic link between HOXB13 and β -catenin/TCF signaling, a recombinant adenovirus expressing HOXB13 (Ad-GFP-HOXB13) was constructed and infected into the aforementioned cells at 10 multiplicity of infection. As shown in Fig. 4A, Ad-GFP-HOXB13 virus showed a greater killing effect on CV-1 cells at day 10 after infection compared with Ad-GFP control virus. However, the same study could not be accomplished in PCa cells, including LNCaP, CWR22RV, PC3, and MDA PCa 2b, because of the viral cytotoxic effect on PCa cells, as reported previously (21). We then studied whether HOXB13 regulates the expression of TCF-4, the most abundant TCF in PCa, or TCF-4-regulated genes. Cells were infected with Ad-GFP-HOXB13 virus, along with Ad-GFP control virus (100 multiplicity of infection). At 48 h after infection, whole cell lysates were prepared and analyzed by Western blot. As demonstrated in Fig. 4B, HOXB13 significantly down-regulated the expression of TCF-4 in CWR22RV and PC3 cells. Alteration of TCF-4 expression in CV-1 cells could not be determined, because antihuman TCF-4

antibodies (Upstate Biotechnology) did not recognize the simian form of TCF-4. The expression of c-myc, a TCF-4 responsive protein, was markedly decreased in most cells. Suppression of cyclin D1 by HOXB13, however, was slight. Presumably this was because the regulation of cyclin D1 requires the long-term expression of HOXB13, rather than a transient expression from a recombinant virus. As predicted in the reporter gene transcription assay shown in Fig. 3D, the level of β -catenin was not affected by HOXB13.

To see the transcription-suppressive effect of HOXB13 in natural promoter, we used *c-myc* promoter spanning a 2.5-kb region containing two copies of TCF-4RE (22). As shown in Fig. 4C, HOXB13 suppressed *c-myc* promoter activity in CWR22RV cells. Next, we investigated whether suppression of TCF-4 by HOXB13 is mediated at the RNA level. After viral infection into PC3 cells, total RNA was extracted, followed by RT-PCR analysis. As shown in Fig. 4D, HOXB13 virus significantly suppressed TCF-4 RNA levels. Attempts to define a mechanistic link between HOXB13 and the promoter region(s) of TCF-4 are in progress. Currently, there is a lack of information on the TCF-4 promoter region (23).

DISCUSSION

Alterations in the level of expression or mutant forms of *HOX* genes may be important in tumorigenesis, *e.g.*, *HOXA9* overexpression transforms myeloid cells, causing leukemia in animal models




Fig. 4. HOXB13 inhibits cell growth by suppressing T-cell factor 4 (*TCF-4*) transcription factor. In **A**, Ad-green fluorescence protein (*GFP*)-*HOXB13* was infected into CV-1 cells, along with control Ad-GFP virus. Cells were monitored by fluorescence microscopy for 10 days. In **B**, cells indicated were infected with either Ad-GFP-*HOXB13* or Ad-GFP. After 48 h, whole cell lysates were prepared, followed by Western blot analysis for the detection of each protein indicated. In **C**, HOXB13 was tested for its effect on reporter construct containing natural *c-myc* promoter (pBV-*cmyc-luc*). In **D**, viruses were infected into PC3 cells as described above, followed by extraction of total RNA. Reverse transcription-PCR was performed to demonstrate the expression of *HOXB13*, *TCF-4*, and β -actin.

(24). HOXB7 is constitutively expressed in melanomas, and transfection of antisense *HOXB7* inhibits cellular proliferation and expression of basic fibroblast growth factor through which HOXB7 directly transactivates (25). Some HOX proteins, including HOXC6 and HOXC8, are involved in PCa progression and overexpressed in more advanced metastatic and recurrent PCa (26, 27).

HOXB13 is the most 5' gene in a cluster on chromosome 17q21, and the mouse and human homologues share 100% amino acid identity in the homeodomain and ~91% in the remainder of the gene (10). Expression of mouse *Hoxb13* is restricted to the terminal spinal cord, hindgut, and urogenital sinus during embryogenesis (10), albeit very low expression of *Hoxb13* is reported in skin (18, 28). In the urogenital sinus, *Hoxb13* starts to express concomitantly with the formation of the prostatic bud in the later stage of embryogenesis. Its expression in adult mouse is confined to the epithelial cells of the prostate, distal colon, and rectum in an androgen-independent fashion (11). *Hoxb13* knockout mice displayed overgrowth in all major structures derived from the tail bud (12) and ventral prostate-specific defects, including defects in epithelial cell morphology and complete loss of secretory proteins (13). These mice showed similar phenotypes to *Nkx3.1* mutant mice and developed a swollen prostate in older mutant mice. However, *Hoxb13* mutant mice did not reveal any phenotypes of prostatic intraepithelial neoplasia, which may require the loss of another regulatory protein, such as *Nkx3.1*. The misexpression of CD44 in luminal epithelial cells observed in these mice is consistent with preneoplastic lesions in many tissue types (29, 30).

The results shown here demonstrate that forced expression of HOXB13 inhibited the growth of PC3 PCa cells. *HOXB13* is reported to be preferentially up-regulated in PCas and PC3 cells (31, 32). Although we have seen that the level of *HOXB13* expression is at least maintained in tumors, we believe that there are subpopulations of both *HOXB13*-positive and -negative PCa cells, which cannot be distinguished by routine RNA analyses. Although Johnson *et al.* (32)

suggested, without showing data, that PC3 cells express HOXB13 and its antisense had an antiproliferative effect on PC3 cells, we have consistently observed that *HOXB13* is not expressed in androgen receptor (AR)-deficient PCa cells (PC3 and DU145), although AR-expressing PCa cells retain *HOXB13* expression (LNCaP, MDA PCa 2b, and LAPC-4).

We have observed that the growth-suppressive effect of HOXB13 was accompanied by a remarkable change in cell morphology in PC3 cells, as shown in Fig. 1B. HOXB13 drove PC3 cells to adopt a gigantic cobblestone appearance, a characteristic phenotype of epithelial cells resulting from terminal differentiation to the mitotically inactive state. This phenomenon has been described to explain the antiproliferative activity of the *Notch* gene in PCa cells (33). HOX proteins are known to be transcription factors. A low level of nuclear HOXB13 was detected in epidermis but not developing dermis, implying that a decrease of HOXB13 is preferred in cells in a differentiated and proliferative mode (18). In addition, *HOXB13* knockout mice exhibited a more fetal-like adult skin, along with enhanced wound healing and higher levels of hyaluronan, considered to inhibit differentiation and promote cell proliferation (17). Altogether, HOXB13 may inhibit cell growth by forcing PC3 cells into a terminal differentiated state. We demonstrated that HOXB13-mediated growth suppression of PC3 cells was accomplished through the down-regulation of TCF-4 expression and subsequently its responsive genes, including *c-myc* and cyclin D1. Suppression of TCF-4 expression was a specific effect by HOXB13; another homeodomain protein, NKX3.1, did not suppress TCF-4-mediated signaling. This result suggests that HOXB13-mediated growth suppression is not caused by nonspecific cell differentiation.

Involvement of HOXB13 in cell cycle arrest was somewhat inconsistent among the transfectants, although the growth-suppressing effect in three of four clones resulted from G₁ arrest. Clone 3 seems to manifest no alteration of cell cycle distribution because of a low level

of HOXB13, although clone 3 grew more slowly than a neo-clone in the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide *in vitro* proliferation assay. It is not unusual for random incorporation of DNA into chromosome to have a growth-regulating effect. A connection between HOX transcription factors and cell cycle progression has been reported, *e.g.*, subgroups, such as HOXA13, HOXC10, and HOXC13, interacted with a sequence within the DNA region of replication associated with the human laminin B2 genes (34). HOX11 abrogated the G₂ cell cycle checkpoint by interacting with protein phosphatase 2A (35).

TCF-4 belongs to a family of transcription factors possessing a high mobility group (HMG) DNA-binding box and facilitates the assembly of multiprotein enhancer complexes. Because TCF-4 is deeply involved in the adenomatous polyposis coli/ β -catenin/TCF pathway, it is thought to be a key cell developmental and growth regulatory protein. In fact, TCF-4 knockout mice die shortly after birth and show an absence of proliferative crypt regions between the villi of the colon (36). TCF-4 has been known to be widely expressed in PCa cells (20). Because TCF-4 is negatively regulated by HOXB13, it is conceivable that tumors acquire a growth advantage from the loss of HOXB13. Although expression of *HOXB13* did not seem to be altered in PCa, >70% of colorectal cancers lost *HOXB13* expression,⁵ suggesting that expression of HOXB13 may be affected by the AR status. In fact, Shen *et al.* (37) have reported the possible involvement of HOXB13 in the regulation of hormone-mediated hormone receptor activation by its interaction with CBP/p300 coactivators. We also observed that HOXB13 is involved in the suppression of androgen-activated AR transcriptional activity.⁶ Therefore, the growth-suppressive function of HOXB13 seems to be accomplished by affecting multiple signaling pathways in prostate tumors. Additional studies are needed to elucidate the association between HOXB13 and AR as a key part of the process of prostate development and tumorigenesis.

REFERENCES

1. Tanaka M, Komuro I, Inagaki H, Jenkins NA, Copeland NG, Izumo S. Nkx3.1, a murine homolog of *Drosophila* bagpipe, regulates epithelial ductal branching and proliferation of the prostate and palatine glands. *Dev Dyn* 2000;219:248–60.
2. Abdulkadir SA, Magee JA, Peters TJ, et al. Conditional loss of Nkx3.1 in adult mice induces prostatic intraepithelial neoplasia. *Mol Cell Biol* 2002;22:1495–503.
3. Bhatia-Gaur R, Donjacour AA, Sciavolino PJ, et al. Roles for Nkx3.1 in prostate development and cancer. *Genes Dev* 1999;13:966–77.
4. Bowen C, Bubendorf L, Voeller HJ, et al. Loss of NKX3.1 expression in human prostate cancers correlates with tumor progression. *Cancer Res* 2000;60:6111–5.
5. Podlasek CA, Clemens JQ, Bushman W. Hoxa-13 gene mutation results in abnormal seminal vesicle and prostate development. *J Urol* 1999;161:1655–61.
6. Podlasek CA, Duboule D, Bushman W. Male accessory sex organ morphogenesis is altered by loss of function of Hoxd-13. *Dev Dyn* 1997;208:454–65.
7. Podlasek CA, Seo RM, Clemens JQ, Ma L, Maas RL, Bushman W. Hoxa-10 deficient male mice exhibit abnormal development of the accessory sex organs. *Dev Dyn* 1999;214:1–12.
8. Prins GS, Birch L, Habermann H, et al. Influence of neonatal estrogens on rat prostate development. *Reprod Fertil Dev* 2001;13:241–52.
9. Warot X, Fromental-Ramain C, Fraulob V, Chambon P, Dolle P. Gene dosage-dependent effects of the Hoxa-13 and Hoxd-13 mutations on morphogenesis of the terminal parts of the digestive and urogenital tracts. *Development* 1997;124:4781–91.
10. Zeltser L, Desplan C, Heintz N. Hoxb-13: a new Hox gene in a distant region of the HOXB cluster maintains colinearity. *Development* 1996;122:2475–84.
11. Sreenath T, Orosz A, Fujita K, Bieberich CJ. Androgen-independent expression of hoxb-13 in the mouse prostate. *Prostate* 1999;41:203–7.
12. Economides KD, Zeltser L, Capecchi MR. Hoxb13 mutations cause overgrowth of caudal spinal cord and tail vertebrae. *Dev Biol* 2003;256:317–30.
13. Economides KD, Capecchi MR. Hoxb13 is required for normal differentiation and secretory function of the ventral prostate. *Development* 2003;130:2061–9.
14. Wu TT, Sikes RA, Cui Q, et al. Establishing human prostate cancer cell xenografts in bone: induction of osteoblastic reaction by prostate-specific antigen-producing tumors in athymic and SCID/bg mice using LNCaP and lineage-derived metastatic sublines. *Int J Cancer* 1998;77:887–94.
15. Lee SJ, Kim HS, Yu R, et al. Novel prostate-specific promoter derived from PSA and PSMA enhancers. *Mol Ther* 2002;6:415–21.
16. Jung C, Ou YC, Yeung F, Frieron HF Jr, Kao C. Osteocalcin is incompletely spliced in non-osseous tissues. *Gene* 2001;271:143–50.
17. Mack JA, Abramson SR, Ben Y, et al. Hoxb13 knockout adult skin exhibits high levels of hyaluronan and enhanced wound healing. *FASEB J* 2003;17:1352–4.
18. Komuves LG, Ma XK, Stelnicki E, Rozenfeld S, Oda Y, Largman C. HOXB13 homeodomain protein is cytoplasmic throughout fetal skin development. *Dev Dyn* 2003;227:192–202.
19. Resnitzky D, Gossen M, Bujard H, Reed SI. Acceleration of the G1/S phase transition by expression of cyclins D1 and E with an inducible system. *Mol Cell Biol* 1994;14:1669–79.
20. Chesire DR, Ewing CM, Gage WR, Isaacs WB. In vitro evidence for complex modes of nuclear beta-catenin signaling during prostate growth and tumorigenesis. *Oncogene* 2002;21:2679–94.
21. Ko SC, Gotoh A, Thalmann GN, et al. Molecular therapy with recombinant p53 adenovirus in an androgen-independent, metastatic human prostate cancer model. *Hum Gene Ther* 1996;7:1683–91.
22. He TC, Sparks AB, Rago C, et al. Identification of c-MYC as a target of the APC pathway. *Science* 1998;281:1509–12.
23. Duval A, Rolland S, Tubacher E, Bui H, Thomas G, Hamelin R. The human T-cell transcription factor-4 gene: structure, extensive characterization of alternative splicing, and mutational analysis in colorectal cancer cell lines. *Cancer Res* 2000;60:3872–9.
24. Golub TR, Slonim DK, Tamayo P, et al. Molecular classification of cancer: class discovery and class prediction by gene expression monitoring. *Science* 1999;286:531–7.
25. Kroon E, Kros J, Thorsteinsdottir U, Baban S, Buchberg AM, Sauvageau G. Hoxa9 transforms primary bone marrow cells through specific collaboration with Meis1a but not Pbx1b. *EMBO J* 1998;17:3714–25.
26. Singh D, Febbo PG, Ross K, et al. Gene expression correlates of clinical prostate cancer behavior. *Cancer Cell* 2002;1:203–9.
27. Waltregny D, Alami Y, Clausse N, de Leval J, Castronovo V. Overexpression of the homeobox gene HOXC8 in human prostate cancer correlates with loss of tumor differentiation. *Prostate* 2002;50:162–9.
28. Stelnicki EJ, Arbeit J, Cass DL, Saner C, Harrison M, Largman C. Modulation of the human homeobox genes PRX-2 and HOXB13 in scarless fetal wounds. *J Invest Dermatol* 1998;111:57–63.
29. Lagorce-Pages C, Paraf F, Dubois S, Belghiti J, Flejou JF. Expression of CD44 in premalignant and malignant Barrett's oesophagus. *Histopathology* 1998;32:7–14.
30. Wimmel A, Kogan E, Ramaswamy A, Schuermann M. Variant expression of CD44 in preneoplastic lesions of the lung. *Cancer* 2001;92:1231–6.
31. Grossmann ME, Wood M, Celis E. Expression, specificity and immunotherapy potential of prostate-associated genes in murine cell lines. *World J Urol* 2001;19:365–70.
32. Johnson PH, Walker RP, Jones SW, et al. Multiplex gene expression analysis for high-throughput drug discovery: screening and analysis of compounds affecting genes overexpressed in cancer cells. *Mol Cancer Ther* 2002;1:1293–304.
33. Shou J, Ross S, Koeppen H, de Sauvage FJ, Gao WQ. Dynamics of notch expression during murine prostate development and tumorigenesis. *Cancer Res* 2001;61:7291–7.
34. de Stanchina E, Gabellini D, Norio P, et al. Selection of homeotic proteins for binding to a human DNA replication origin. *J Mol Biol* 2000;299:667–80.
35. Kawabe T, Muslin AJ, Korsmeyer SJ. HOX11 interacts with protein phosphatases PP2A and PP1 and disrupts a G2/M cell-cycle checkpoint. *Nature* 1997;385:454–8.
36. Korinek V, Barker N, Moerer P, et al. Depletion of epithelial stem-cell compartments in the small intestine of mice lacking Tcf-4. *Nat Genet* 1998;19:379–83.
37. Shen WF, Krishnan K, Lawrence HJ, Largman C. The HOX homeodomain proteins block CBP histone acetyltransferase activity. *Mol Cell Biol* 2001;21:7509–22.

⁵ C. Jung, S.J. Lee, L. Cheng, T. A. Gardner, M.H. Jeng, and C. Kao. HOXB13 functions as a tumor suppressor in colorectal cancer, manuscript in preparation.

⁶ C. Jung, R. S. Kim, H.J. Zhang, S.J. Lee, and M.H. Jeng. HOXB13 induces growth suppression of prostate cancer cells as a repressor of hormone-activated androgen receptor signaling, submitted for publication.