

MOK™

■ *Complete* Media Optimization Kit™

Cat. No. 0185

The *Complete* Media Optimization Kit™ determines the best medium formulation for maximizing accumulation of recombinant proteins expressed in *E. coli*, utilizing a series of Athena's superior proprietary expression media as well as two reference media.



Athena Enzyme Systems™

888-892-8408

www.athenaes.com

Application Manual

V 4.0

MOK™

Complete Media Optimization Kit™

Cat. No. 0185

Application Manual

V. 4.0

© Athena Environmental Sciences, Inc.

Turbo Broth™, Superior Broth™, Power Broth™, Hyper Broth™, LB*Booster™, Augmedium™, and MOK™ (Media Optimization Kit™) are all trademarks of Athena Environmental Sciences, Inc. Athena Enzyme Systems™ is a division and trademark of Athena Environmental Sciences, Inc. AthenaES™ is a trademark of Athena Environmental Sciences, Inc. and the AthenaES™ logo is a registered logo of Athena Environmental Sciences, Inc.



Athena Enzyme Systems™

1450 South Rolling Road

Baltimore, MD 21227

USA

T (MD): 410-455-6319

T (USA): 888-892-8408

F: 410-455-1155

aesinfo@athenaes.com

a division of Athena Environmental Sciences, Inc.

Complete MOK™: Media Optimization Kit™

Application Manual

Table of Contents

Introduction	4
Principle of the Kit	4
Kit Components	5
Protocols:	
Preparation: Dry Powder Media	5
Media Screening Protocol	6
Tips	8
Product Warranty	9
Product Use Limitations	9
Ordering Information	10

Choosing the right culture conditions can be as important as choosing the right expression system to produce your protein.

Introduction

The Complete Media Optimization Kit™ was designed to provide the researcher with a tool for determining the best available medium formulation for the production of recombinant proteins in *Escherichia coli*. The level of expression of a given protein is dependent upon the composition of the medium used. In the course of optimizing the production of recombinant proteins for clients, Athena's scientists have developed several media that were specifically formulated to maximize the accumulation of recombinant proteins expressed in *E. coli*. Four of these unique blends, Turbo Broth™, Superior Broth™, Power Broth™, Hyper Broth™, including their APF counterparts, are contained in the Complete Media Optimization Kit™ along with reference media LB Broth and Glucose M9Y. These blends of media have proven to be the most widely applicable formulations showing consistently superior performance over the traditional medium, LB Broth.

Historically, *E. coli* has been cultivated in LB Broth¹ and many gene expression protocols recommend this medium.² It should be pointed out that this medium was developed in the 1950's, nearly 20 years before the first gene was cloned and 30 years before recombinant protein expression became routine. Therefore, while LB Broth has proved to be very useful for cultivating *E. coli*, it was not specifically designed with the intention of maximizing the expression of recombinant proteins. Moreover, LB Broth is not normally supplemented with a carbon source nor is it buffered. Thus, the growth yields that one can obtain with LB are limited. Moreover, we have found that many proteins are not readily expressed in LB (See Figure 1. on page 7 for an example).

Principle of the Kit

The principal purpose of the optimization kit is to allow for rapid identification of a suitable medium without the need for extensive optimization testing or development work. The test simply involves expressing the desired protein using each of the media provided in the kit. The relative level of expression of the target protein, after induction of expression, is determined by SDS-PAGE using a qualitative assessment. In some cases an immunoblot or functional assay may be appropriate.

The optimum conditions for the expression of a recombinant protein requires attention to four culture-related parameters. These include the strain employed, the medium composition, the time-course of induction, and the concentration of inducing agent. We recommend performing the culture optimization tests in the order listed. Thus, the Complete Media Optimization Kit™ should be used after selecting the best available strain.

Kit Components

Individually packaged, ready-to-use powdered media to make 1 liter of each broth.

Complete Media Optimization Kit™			
Component	Catalog Number	Component	Catalog Number
LB Broth (Miller)	0103	APF LB (Miller)*	0133
Turbo Broth™	0104	Turbo Prime™*	0110
Superior Broth™	0105	Superior Prime™*	0111
Power Broth™	0106	Power Prime™*	0112
Hyper Broth™	0107	Glucose M9Y*	0108
Turbo Prime-olate Broth™	0160	Superior Prime-olate™*	0161
Power Prime-olate Broth™	0162	Glucose Nutrient Mix	0109

*APF formulations

Protocols

Preparation: Dry Powder Media

1. Dissolve the contents of each of the media packets in deionized water as directed on the individual packets.
2. Add 4 mL of glycerol to the Turbo Broth™ and Power Broth™, Turbo Prime Broth™ and Power Prime Broth™, or to the Turbo Prime-olate Broth™ and Power Prime-olate Broth™ solutions.
3. Dispense desired volume into appropriate bottles or flasks. (We recommend 2 x 500mL glass bottles.)
4. Autoclave at 121°C for 20 min. The autoclaved media without antibiotics are stable for 6 months at 4°C.
5. Dissolve the contents of the Glucose Nutrient Mix in 100 mL deionized water and filter sterilize using a 0.2µm filter.
6. Add 50 mL of the sterile Glucose Nutrient Mix to 1 liter of Hyper Broth™ and 20 mL to 1 liter of Glucose M9Y using aseptic technique.
7. Add sterile antibiotics as needed.

Media Screening Protocol

1. Materials

- 1.1. 25 mL of each culture medium in 250 mL baffled bottomed flasks.
- 1.2. Wash Buffer: 40 mM sodium phosphate pH 7.5, 150 mM NaCl
- 1.3. 2x SDS-PAGE Loading Dye: 125 mM Tris-Cl pH 6.8, 4% SDS (w/v), 0.005% bromophenol blue (w/v), 20% glycerol (v/v), 5% β-mercaptoethanol (v/v)
- 1.4. Tris-Glycine SDS-polyacrylamide gel of appropriate composition.

2. Methods

- 2.1. Inoculate a single colony of the recombinant strain into 10 mL of LB Broth in a baffled flask. Incubate at 37° C overnight.
- 2.2. Inoculate 25 mL of each media with 1 mL of the overnight culture. Incubate the cultures at 37° C until the OD₆₀₀ reaches 0.6.
- 2.3. Remove a 1 mL sample ("pre-induction"), harvest the cells in a pre-weighed microfuge tube, and store on ice.
- 2.4. Add inducer (see Tip 1) and continue incubating for 3 hours (see Tip 2).
- 2.5. Remove a 1 mL sample ("post-induction") and process as in step 2.3.
- 2.6. Harvest the remainder of the culture, wash with 10 mL of wash buffer, determine the mass of the cell pellet, and store the cell pellets at -80° C. (see Tip 3.)
- 2.7. Analyze for expression of the target protein as follows:
 - 2.7.1. To determine protein production per mL of culture:
 - 2.7.1.1. Suspend the cell pellets from the pre- and post-induction samples in 0.5 mL of water.
 - 2.7.1.2. Mix 5 µL of each cell suspension with 7.5µL water and 12.5µL 2x SDS-PAGE loading buffer. Heat at 100° C for 5 minutes and load 10 µL per lane of acrylamide gel.
 - 2.7.2. To determine the relative level of expression:
 - 2.7.2.1. Suspend the cell pellets from the pre- and post-induction samples in water to a density of 10 OD/mL

2.7.2.2. Mix 5 μL of each cell suspension with 7.5 μL water and 12.5 μL 2x SDS-PAGE loading buffer. Heat at 100° C for 5 minutes and load 10 μL per lane of acrylamide gel.

2.7.3. Stain the gel with Coomassie Blue, colloidal Coomassie Blue or Silver stain. (see *Tips 4 and 5*).

3. Interpretation

- 3.1. After staining the gel, observe each lane and compare the “pre-induction” sample with the “post-induction” sample from each medium. Elevated expression is indicated by the presence of a unique polypeptide band corresponding to the molecular mass of the target protein in the “post-induction” sample.
- 3.2. Compare the level of target protein from cells grown in each of the six media. Select the medium which produces the highest level of target protein per mL of culture. Figure 1 shows the results of a media screening experiment.
- 3.3. If two or more media give the same level of production per mL, then use the analysis of 2.7.2 to select the medium with the highest relative level of expression.




Figure 1. Media screen analysis by SDS-PAGE of a recombinant protein expressed in *E. coli*. The arrow denotes the recombinant protein. The protein was expressed in strain M15 grown in LB Broth, Glucose M9Y, Hyper Broth™, Power Broth™, Superior Broth™, and Turbo Broth™, lanes 1 to 6, respectively. Lane 7 is Novex SeeBlue™ marker proteins. For this particular protein, Hyper Broth™ was selected for production.

Tips of the Trade

Tip 1: Media Optimization Inducer

The inducer used will depend on the expression system employed. The concentration of inducer is strain-dependent and the optimum concentration should be determined empirically. For *lacP*-based expression systems, 1 mM IPTG is good for the media optimization.

Tip 2: Inclusion Bodies

Some recombinant proteins are expressed in *E. coli* as insoluble particles known as inclusion bodies. The formation of inclusion bodies can not be predicted, but are indicated by the presence of intracellular refractive objects when viewed under oil immersion microscopy.³ The formation of inclusion bodies will not affect the results of the media screen, because the analysis is done on whole cell extracts prepared by boiling the cells in sodium dodecylsulfate. This procedure completely denatures inclusion bodies as well as membrane and cytoplasmic proteins. (In some instances lowering the temperature after induction can increase the amount of soluble protein recovered. This should be determined experimentally.)

Tip 3: Cell Paste

Once the medium yielding the highest level of expression has been determined, the cell paste can be used to prepare a small-scale extract.

Tip 4: SDS-PAGE gels

Coomassie Blue stain should be sufficient to visualize the expression of a recombinant protein. Silver stain, while allowing detection of smaller amounts of protein, is more difficult to interpret and should only be used for examining whole cell extracts which are separated by SDS-polyacrylamide gels that are 20 cm in length or longer. The long gel will give better resolution of individual polypeptide bands.

Tip 5: Alternative Techniques

Alternative techniques can be applied to the media screen analysis. Immunoblot or functional assays can be employed as appropriate. Care should be taken when using functional assays by first demonstrating that there is no interfering activity contributed by the host. In most cases, the SDS-PAGE analysis is the method of choice during the early stages of developing the expression system. Immunoblots should be used when the Coomassie blue stain does not reveal any expressed protein.

Technical Assistance

The scientific staff of the Athena Enzyme Systems™ are specialists in the expression of recombinant proteins in microbial systems. They have extensive expertise in all aspects of protein expression from the construction of expression vectors to the commercial production of recombinant proteins. No matter what your question, please feel free to ask them for help. A technical support scientist can be reached at support@athenaes.com.

Product Use Limitations

The Complete Media Optimization Kit was designed and is sold for research use only. It should not be used for human diagnosis or drug use or administered to humans unless expressly cleared for that purpose by the appropriate regulatory authorities in the country of use. All due care and attention should be exercised in the handling of the materials contained in the kit.

Product Warranty

AthenaES™ guarantees the quality and performance of the media and reagents contained in this kit for the cultivation of *E. coli*. The suitability of a medium formulation or additive for a particular use is the responsibility of the end user. No guarantee is made that a given protein will be expressed when applying this kit. AthenaES™ will replace the product free of charge if it does not conform to the stated specifications. Notice for replacement must be received within 60 days of opening the product.

References

1. Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. Molecular cloning: a laboratory manual, 2nd edition. Cold Spring Harbor Laboratory, Cold Spring Harbor, New York
2. Luria, S. E., and J. W. Burrous. 1955. Hybridization between *Escherichia coli* and *Shigella*. J. Bacteriol. 74:461-476.
3. Broedel, Jr., S. E., S. M. Papciak, and W. R. Jones. 2001. The selection of optimum medium formulations for improved expression of recombinant proteins in *E. coli*. Athena Enzyme Systems Technical Bulletin, January 2001.

Ordering Information

To place an order:

Phone: 1-888-892-8408 Email: orders@athenaes.com
 Fax: 410-455-1155 Website: www.athenaes.com

Or visit our website to order through one of our international distributors.

When placing an order, please provide the following:

- Institution name and customer service account
- Purchase order number
- Catalog number(s) or names of products and quantity of item(s)
- Billing and shipping address
- Contact name and telephone number

Delivery:

Telephone orders received Monday through Friday before 12 noon will be shipped that day. All other orders will be shipped the next business day, unless otherwise stipulated.

Complete Media Optimization Kit™ Product Ordering Information		
Catalog Number	Product	Size
0103	LB Miller	500 g
0133	APF LB (Miller)*	500 g
0104	Turbo Broth™	500 g
0110	Turbo Prime™*	500 g
0160	Turbo Prime-olate™*	500 g
0105	Superior Broth™	500 g
0111	Superior Prime™*	500 g
0161	Superior Prime-olate™*	500 g
0106	Power Broth™	500 g
0112	Power Prime™*	500 g
0162	Power Prime-olate™*	500 g
0107	Hyper Broth™*	500 g
0108	Glucose M9Y*	500 g
0109	Glucose Nutrient Mix	150 g
Companion Products		
Catalog Number	Product	Size
0123 (0124)	Augmedium™*	100 ml (500 ml)
0125 (0126)	LB Booster™*	100 ml (500 ml)
New! Ready-to-use Sterile Liquid Kit		
Catalog Number	Product	Size
0187	Expression Media Selection Kit™	15 ml per media (fifteen formulations)

*APF formulations